

A PHOTOMETRIC VARIABILITY SURVEY OF FIELD K AND M DWARF STARS WITH HATNET

J. D. HARTMAN¹, G. Á. BAKOS¹, R. W. NOYES¹, B. SÍPŐCZ^{1,2}, G. KOVÁCS³, T. MAZE⁴, A. SHPORER⁴, A. PÁL^{1,2}*Draft version July 16, 2009*

ABSTRACT

Using light curves from the HATNet survey for transiting extrasolar planets we investigate the optical broad-band photometric variability of a sample of 27,560 field K and M dwarfs selected by color and proper-motion ($V - K \gtrsim 3.0$, $\mu > 30$ mas/yr, plus additional cuts in $J - H$ vs. $H - K_S$ and on the reduced proper motion). We apply a variety of variability selection algorithms on the light curves to search for periodic and quasi-periodic variations, and for large-amplitude, long-duration flare events. To set the selection thresholds we conduct Monte Carlo simulations of light curves with realistic noise properties. A total of 3496 stars exhibit potential variability, including 95 stars with eclipses and 60 stars with flares. Based on a visual inspection of these light curves and an automated blending classification, we select 1928 stars, including 79 eclipsing binaries, as secure variable star detections that are not high probability blends. We find that only 43 of these stars, including 7 of the eclipsing binaries, have previously been identified as variables or are blended with previously identified variables. One of the newly identified eclipsing binaries is 1RXS J154727.5+450803, a known $P = 3.55$ day, late M-dwarf SB2 system, for which we derive preliminary estimates for the component masses and radii of $M_1 = M_2 = 0.258 \pm 0.008 M_\odot$ and $R_1 = R_2 = 0.289 \pm 0.007 R_\odot$. The radii of the component stars are larger than theoretical expectations if the system is older than ~ 200 Myr. The majority of the variables are heavily spotted BY Dra-type stars for which we determine rotation periods. This is the largest sample of photometric rotation periods for field K and M dwarfs published to date. Using this sample, we investigate the relations between period, color, age, and activity measures, including optical flaring, for K and M dwarfs. We find that the fraction of stars that are variable with peak-to-peak amplitudes greater than 0.01 mag increases exponentially with the $V - K_S$ color such that approximately half of field dwarfs in the solar neighborhood with $M \lesssim 0.2 M_\odot$ are variable at this level. Our data hints at a change in the rotation-activity-age connection for stars with $M \lesssim 0.25 M_\odot$.

Subject headings: stars: late-type, low-mass, rotation, flare, fundamental parameters (masses, radii)
— binaries: eclipsing — surveys — catalogs — techniques: photometric

1. INTRODUCTION

With the advent of large-scale automated time-series photometric surveys to search for microlensing events, transiting planets, and supernovae, well-sampled light curves for millions of stars have been, and are continuing to be, collected. While the primary scientific goals of these surveys are to search for rare phenomena, the enormous databases of light curves that are a by-product of these surveys presents a great opportunity to study many other topics related to stellar variability (e.g. Paczyński 1997). With some exceptions (e.g. Hartman et al. 2004; Creevey et al. 2005; Parley et al. 2006; Norton et al. 2007; Beatty et al. 2007; Karoff et al. 2007; Shporer et al. 2007; Fernandez 2009) the copious data produced by dedicated wide-field transit surveys has been relatively under-utilized for studying topics unrelated to transiting planets. In this paper we use data from the Hungarian-made Automated Telescope Network (HATNet; Bakos et al. 2004) project to study the variability of probable K and M dwarf field stars.

Combining photometric observations with proper motion measurements is an effective method for selecting

nearby dwarf stars. This technique is routinely used in the search for cool stars in the solar neighborhood (e.g. Reyle & Robin 2004), and has been suggested as an effective method for screening giants from transit surveys (Gould & Morgan 2003). By selecting red, high proper motion stars, it is possible to obtain a sample that consists predominately of nearby K and M-dwarfs, with very few luminous, distant giants. Whereas a general variability survey of Galactic field stars will yield a mélange of objects that are often difficult to classify without detailed follow-up (e.g. Hartman et al. 2004; Shporer et al. 2007), by focusing a survey on red, high proper motion stars, one can narrow in on a few specific topics related to low-mass stars.

Main-sequence stars smaller than the Sun are not known to exhibit significant pulsational instabilities, but they may exhibit the following types of photometric variability: 1. Variability due to binarity (either eclipses or proximity effects such as ellipsoidal variability). 2. Variability due to the rotational modulation or temporal evolution of starspots. 3. Flares. We discuss each type of variability, and what might be learned from studying it, below.

1.1. Low-Mass Eclipsing Binaries

In recent years there has been an increasing number of EBs discovered with K and M dwarf components (see the list of 13 such binaries compiled by Devor et al. 2008,

¹ Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138, USA

² Department of Astronomy, Eötvös Loránd University, Budapest, Hungary.

³ Konkoly Observatory, Budapest, Hungary.

⁴ Wise Observatory, Tel Aviv University, Tel Aviv, Israel

note that most of these were found in the last 5 years). From these discoveries it has become clear that the radii of early M dwarfs and late K dwarfs are somewhat larger than predicted by theory (the number typically stated is 10%; Torres & Ribas 2002; Ribas 2003; López-Morales 2005; Ribas 2006; Beatty et al. 2007; Fernandez 2009).

Most of the binaries found to date have periods shorter than a few days and are expected to have rotation periods that are tidally synchronized to the orbital period. The rapid rotation in turn yields enhanced magnetic activity on these stars compared to isolated, slowly rotating stars. It has been suggested that the discrepancy between theory and observation for these binary star components may be due to their enhanced magnetic activity inhibiting convection. (Ribas 2006; Torres et al. 2006; López-Morales 2007; Chabrier et al. 2007). Support for this hypothesis comes from interferometric measurements of the luminosity-radius relation for inactive single K and M dwarfs which appears to be in agreement with theoretical predictions (Demory et al. 2009). There is also evidence that the discrepancy may be correlated with metallicity (López-Morales 2007). Testing these hypotheses will require finding additional binaries spanning a range of parameters (mass, rotation/orbital period, metallicity, etc.).

As transiting planets are discovered around smaller stars, the need for models that provide accurate masses and radii for these stars has become acute. For example, the errors in the planetary parameters for the transiting Super-Neptune GJ 436b appear to be dominated by the uncertainties in the stellar parameters of the $0.45 M_{\odot}$ M-dwarf host star (Torres 2007, note that the author gives errors in the mass and radius for the star of $\sim 3\%$ assuming that the theoretical models for the luminosity of M-dwarfs are accurate while making no such assumption about the radius). Having a large sample of low mass stars with measured masses and radii would enable the determination of precise empirical relations between the parameters for these stars.

1.2. Stellar Rotation

The rotation period is a fundamental measurable property of a star. For F, G, K and early M main sequence stars there is a well-established relation between rotation, magnetic activity and age (e.g. Skumanich 1972; Noyes et al. 1984; Pizzolato et al. 2003). In addition to illuminating aspects of stellar physics, this relation in practice provides the best method for measuring the ages of field main sequence stars (Mamajek & Hillenbrand 2008; Barnes 2007).

For late M-dwarfs the picture is less clear. From a theoretical standpoint one might expect that fully convective stars should not exhibit a rotation-activity connection if the connection is due to the $\alpha\Omega$ -dynamo process, which operates at the interface of the radiative and convective zones in a star, and is thought to generate the large scale magnetic fields in the Sun (see for example the discussion by Mohanty & Basri 2003). Nonetheless, several studies have found evidence that the rotation-activity connection continues to very late M-dwarfs (Delfosse et al. 1998; Mohanty & Basri 2003; Reiners & Basri 2007). In these studies the rotation period is inferred from the projected rotation velocity $v \sin i$, which is measured spectroscopically, while the degree of magnetic activity is es-

timated by measuring either the $H\alpha$ emission or X-ray emission. Rotation studies of this sort suffer both from the inclination axis ambiguity, and from low sensitivity to slow rotation. In practice it is very difficult to measure $v \sin i$ values less than ~ 1 km/s, which generally means that it is only possible to place lower limits on the period for late M-dwarf stars with periods longer than ~ 10 days. Moreover, because low-mass stars are intrinsically faint, these studies require large telescopes to obtain high-resolution, high S/N spectra, so that typically only a few tens of stars are studied at a time.

There are two techniques that have been used to directly measure stellar rotation periods. The first technique, pioneered by Wilson & Bappu (1957), is to monitor the emission from the cores of the CaII H and K lines, searching for periodic variations. The venerable Mount-Wilson Observatory HK project has used this technique to measure the rotation periods of more than 100 slowly rotating dwarfs and giant stars (Wilson 1978; Duncan et al. 1991; Baliunas et al. 1995, 1996). Alternatively, if a star has significant spot-coverage it may be possible to measure its rotation period by detecting quasi-periodic variations in its broad-band photometric brightness. Studies of this sort have been carried out in abundance for open clusters (e.g. Radick et al. 1987) as well as for some field stars (e.g. Strassmeier et al. 2000). While there are rich samples of rotation periods for K and M dwarfs in open clusters with ages $\lesssim 600$ Myr (Irwin et al. 2006, 2007, 2009a; Meibom et al. 2009; Hartman et al. 2009), the data for older K and M dwarfs is quite sparse. As such, there are few observational constraints on the rotational evolution of these stars after ~ 0.5 Gyr.

Unlike spectroscopic studies, photometric surveys may yield rotation periods for hundreds of stars at a time. There are, however, some drawbacks to these surveys. The spot distribution on the surface of a star may in general be quite complex, so the resulting signal in the light curve will not always take a simple form. Since the number of brightness minima per cycle is not known a priori, there is a risk that the true rotation period may be a harmonic of the measured period. Spots on the Sun come and go on time-scales shorter than the Solar rotation period, and indeed stellar light curves also exhibit secular trends. For long period stars the measured variation time-scale may actually correspond to a spot evolution time-scale rather than the rotation period of the star. For short period stars there may be difficulties in distinguishing spot modulation from binarity effects (though for these stars the rotation period is expected to be tidally synchronized to the orbital period).

Despite these caveats, given the existing uncertainties in the rotation-activity connection for low mass stars and the potential to use rotation as a proxy for age, a large, homogeneously collected sample of photometric rotation periods for field K and M dwarfs could potentially be of high value.

1.3. Flares

Flaring is known to be a common phenomenon among K and M dwarfs. Studies of open cluster and field flare stars have shown that the frequency of flaring increases with decreasing stellar mass, and decreases with increasing stellar age (e.g. Ambartsumyan et al.

1970; Mirzoyan et al. 1989). Significant flaring on these low-mass dwarfs is likely to impact the habitability of any planets they may harbor (e.g. Kasting et al. 1993; Lammer et al. 2007; Guinan & Engle 2009), so determining the frequency of flares, and its connection with other stellar properties such as rotation, has important implications for the study of exoplanets.

1.4. The HATNet Survey

To address these topics we use data from HATNet to conduct a variability survey of K and M dwarfs. The ongoing HATNet project is a wide-field search for transiting extrasolar planets (TEPs) orbiting relatively bright stars. The project employs a network of 7 robotic telescopes (4 in Arizona at Fred Lawrence Whipple Observatory, 2 in Hawaii on the roof of the Sub-Millimeter Array at Mauna Kea Observatory, and 1 in Israel at Wise Observatory; the latter is referred to as WHAT, see Shporer et al. 2009) which have been used to obtain some $\sim 700,000$ images covering approximately 10% of the sky. The survey has generated light curves for approximately 2.5 million stars, from which ~ 900 candidate TEPs have been identified. To date, the survey has announced the discovery of 12 TEPs, including HAT-P-11b (Bakos et al. 2009), a Super-Neptune ($0.08 M_J$) planet that is the smallest found so far by a ground-based transit survey. While the primary focus of the HATNet project has been the discovery of TEPs, some results not related to planets have also been presented. This includes the discovery and analysis of a low-mass M dwarf in a single-lined eclipsing binary (EB) system (Beatty et al. 2007), searches for variable stars in two HATNet/WHAT fields (Hartman et al. 2004; Shporer et al. 2007).

1.5. Overview of the Paper

The structure of the paper is as follows. In § 2 we describe both the HATNet photometric data, and select the sample of field K and M dwarfs. In § 3 we discuss our methods for selecting variable stars. We estimate the degree of blending for potential variables in § 4. We match our catalog of variables to other catalogs in § 5. We discuss the properties of the variables in § 6 including an analysis of one of the EB systems found in the survey. We conclude in § 7. In appendix A we describe the Monte Carlo simulations used in establishing our variability selection thresholds, while in appendix B we present the catalog of variable stars.

2. OBSERVATIONAL DATA

2.1. HATNet Data

The HATNet project, which has been in operation since 2003, uses a network of 7 small (11 cm aperture), autonomous telescopes to obtain time-series observations of stars. For details on the system design see Bakos et al. (2004); here we briefly review a few points that are relevant to the survey presented in this paper. Prior to 2007 September each telescope employed a $2K \times 2K$ CCD and a Cousins I_C filter (Cousins 1976). The $2K \times 2K$ CCDs covered an $8.2^\circ \times 8.2^\circ$ field of view (FOV) at a pixel scale of $14''$. With these CCDs, stars with $7.5 \lesssim I_C \lesssim 14.0$ were observed with a typical per-image photometric precision of a few mmag at the bright end, 0.01 mag at $I_C \sim 11$, and 0.1 mag at $I_C \sim 13.5$. After this date the telescopes

were refitted with $4K \times 4K$ CCDs and Cousins R_C filters. The new CCDs cover a $10.6^\circ \times 10.6^\circ$ FOV at a pixel scale of $9''$. With these CCDs, stars with $8.0 \lesssim R_C \lesssim 15.0$ are observed with a typical per-image photometric precision of a few mmag at the bright end, 0.01 mag at $R_C \sim 12$, and 0.1 mag at $R_C \sim 15$. The exact magnitude limits and precision as a function of magnitude vary within a field due to vignetting, and from field to field due to differences in the reduction procedure used and in the degree of stellar crowding. In 2008 September the filters were changed to Sloan r , though we do not include any observations taken through the new filters in the survey presented here. For both CCD formats, the typical full width at half maximum (FWHM) of the point spread function (PSF) is ~ 2 pixels (i.e. $\sim 30''$ for the $2K$ fields and $\sim 20''$ for the $4K$ fields).

The data for this survey comes from 72 HATNet fields with declinations between $+15^\circ$ and $+52^\circ$. These fields are defined by dividing the sky into $838 7.5^\circ \times 7.5^\circ$ tiles. The survey covers approximately 4000 square degrees, or roughly 10% of the sky.

The data reduction pipeline has evolved over time, as such the fields studied in this survey have not all been reduced in a uniform manner. For simplicity we choose to use the available light curves as is, rather than re-reducing the fields in a consistent manner optimized for finding variable stars rather than TEPs. Both aperture photometry (AP) and image subtraction photometry (ISM) have been used for reductions. Both pipelines were developed from scratch for HATNet. See Pál (2009) for detailed descriptions of both methods.

For both pipelines the Two Micron All-Sky Survey (2MASS; Skrutskie et al. 2006) is used as the astrometric reference. The astrometric solutions for the images are determined using the methods described by Pál & Bakos (2006) and Pál (2009). Photometry is performed at the positions of 2MASS sources transformed to the image coordinate system. For each resulting light curve the median magnitude is fixed to the I_C or R_C magnitude of the source based on a transformation from the 2MASS J, H and K_S magnitudes. For the ISM reduction this magnitude is also used as the reference magnitude for each source in converting differential flux measurements into magnitudes.

Both the AP and ISM pipelines produce light curves that are calibrated against ensemble variations in the flux scale (for AP this is done as a step in the pipeline, for ISM this is an automatic result of the method). For each source, light curves are obtained using three separate apertures. The set of apertures used has changed over time; the most recent reductions use aperture radii of 1.45, 1.95 and 2.35 pixels. Following the post-processing routines discussed below, we adopt a single “best” aperture for each light curve.

The calibrated light curves for each aperture are passed through two routines that remove systematic variations from the light curves that are not corrected in calibrating the ensemble. The first routine (EPD) decorrelates each light curve against a set of external parameters including parameters describing the shape of the PSF, the sub-pixel position of the star on the image, the zenith angle, the hour angle, the local sky background, and the variance of the background (see Bakos et al. 2009). After applying EPD, the light curves

are then processed with the Trend-Filtering Algorithm (TFA; Kovács, Bakos & Noyes 2005) which decorrelates each light curve against a representative sample of other light curves from the field. The number of template light curves used differs between the fields, typically the number is $\sim 8\%$ of the total number of images for that field. In applying the TFA routine we also perform σ -clipping on the light curves since this generally reduces the number of false alarms when searching for transits. For the remainder of the paper we will refer to light curves that have been processed through EPD only, without application of TFA, as EPD light curves, and will refer to light curves that have been processed through both EPD and TFA as TFA light curves. We note that for some fields the EPD light curves were not stored and only TFA light curves are available.

Both of these algorithms tend to improve the signal to noise ratio of transit signals in the light curves, but they may distort the light curves of stars that show large-amplitude, long-period, continuous variability. Additionally the decorrelation against the zenith and hour angles in the EPD routine will tend to filter out real variable star signals with periods very close to a sidereal day or an integer multiple of a sidereal day. The TFA routine in particular may distort long-period signals while increasing the signal to noise ratio of short-period signals. For this reason we analyze both the EPD and TFA light curves, when available, to select variable stars (§ 3). We note that for the analysis in this paper we do not use the signal-reconstruction mode TFA presented by Kovács, Bakos & Noyes (2005). Once a signal is detected, TFA can be run in this mode to obtain a trend-filtered light curve that is free of signal distortions, however for signal detection one must use general TFA since the signal is not known a priori.

Finally, an optimal aperture is chosen for each star. For stars fainter than a fixed limit the smallest aperture is used (to minimize the sky noise), for brighter stars the aperture with the smallest root-mean-square (RMS) light curve is used.

2.2. Composite Light Curves

Because the separation between the HATNet field centers is smaller than the FOV of the HATNet telescopes for both the 2K and 4K CCDs, some stars are observed in multiple fields. These stars may have more than one light curve, which we combine into composite light curves. In making a composite light curve we subtract the median magnitude from each component light curve. For fields reduced with both ISM and AP we use the light curve with the lowest RMS, and in the case of equal RMS we use the ISM light curve. Note that the composite light curve for a star may include a mix of I_C and R_C photometry. While the amplitude of variability may be different from filter to filter, the period and phasing for variations due to eclipses or the rotational modulation of starspots will be independent of bandpass. For simplicity we do not allow for independent amplitudes of different filters in searching for variability using the methods described in § 3; we do not expect this to make a significant difference to period detections. However, we note that a simple merging of photometry from different filters may result in spurious side lobes in the power spectrum; for a more detailed analysis of individual objects this effect

should be considered.

2.3. Selection of the K and M Dwarf Sample

To select the sample of stars that are probable K and M dwarfs we apply cuts on the proper motion and on the color. Proper motion measurements are taken from the PPM-Extended catalog (PPMX; Röser et al. 2008) which provides proper motions with precisions ranging from 2 mas/yr to 10 mas/yr for 18 million stars over the full sky down to a limiting magnitude of $V \sim 15.2$. The PPMX catalog provides complete coverage of the HATNet survey for stars with $V - I_C \lesssim 1.2$ for the 2K fields and for stars with $V - R \lesssim 0.7$ for the 4K fields. For stars redder than these limits, the faint limit of HATNet is deeper than the faint limit of PPMX. We select all stars from this catalog with a proper motion $\mu > 30$ mas/yr. For the color selection we use the 2MASS JHK_S photometry and, where available, V -band photometry from the PPMX catalog, which is taken from the Tycho-2 catalog (Høg et al. 2000) and transformed to the Johnson system by Kharchenko (2001). Only $\sim 4\%$ of the stars have V photometry given in the PPMX catalog, for the majority of stars that do not have V photometry we calculate an approximate V magnitude using

$$V = -0.0053 + 3.5326J + 1.3141H - 3.8331K_S. \quad (1)$$

which is determined from 590 Landolt Standard stars (Landolt 1992) with 2MASS photometry, and is used internally by the HATNet project to estimate the V magnitudes of transit candidates for follow-up observations. We then select stars with $V - K_S > 3.0$ which corresponds roughly to stars with spectral types later than K6 (Bessell & Brett 1988). This selects a total of 471,970 stars from the PPMX catalog, of which 33,177 fall in a reduced HATNet field; of these 32,831 have a HATNet light curve containing more than 1000 points.

Note that extrapolating a V magnitude from near-infrared photometry is not generally reliable to more than a few tenths of a magnitude. We therefore also obtained V magnitudes for stars in our sample by matching to the USNO-B1.0 catalog (Monet et al. 2003). This matching was done after the variability search described in § 3; we choose not to redo the sample selection and the subsequent variability selection. Note that low-mass sub-dwarfs, which have anomalously blue $V - K_S$ values, will pass a selection on $V - K_S > 3.0$ computed using eq. 1 to extrapolate the V magnitude, while they would not necessarily pass a selection using the measured value of $V - K_S$. To transform from the photographic B_U , R_U magnitudes in the USNO-B1.0 catalog to the V -band we use a relation of the form:

$$V = aB_U + bB_U^2 + cR_U + dR_U^2 + eB_U R_U + f \quad (2)$$

with coefficients given separately in table 1 for the $(B_{U,1}, R_{U,1})$, $(B_{U,1}, R_{U,2})$, $(B_{U,2}, R_{U,1})$ and $(B_{U,2}, R_{U,2})$ combinations. These transformations were determined using ~ 1100 stars with both V photometry in the PPMX catalog and USNO-B1.0 (B_U, R_U) photometry. Based on the root-mean-square (RMS) scatter of the post-transformation residuals, we used the $(B_{U,2}, R_{U,2})$, $(B_{U,1}, R_{U,2})$, $(B_{U,2}, R_{U,1})$ and $(B_{U,1}, R_{U,1})$ transformations in order of preference. For stars with neither PPMX V photometry nor USNO-B1.0 photometry, we

used eq. 1. For the remainder of the analysis in this paper, the V magnitude is taken from PPMX (Tycho-2) for 3.1% of the stars in our sample, from USNO-B1.0 for 93.6% of the stars, and is transformed from the 2MASS magnitudes for 3.3% of the stars.

Figure 1 shows the $J - H$ vs. $H - K_S$ color-color diagram for the selected sample. We also show the expected relations for dwarf stars and for giants. The relation for dwarfs is taken from a combination of the Baraffe et al. (1998) 1.0 Gyr isochrone for solar metallicity stars with $0.15 M_\odot \leq M \leq 0.7 M_\odot$ and the Chabrier et al. (2000) models for objects with $M \leq 0.075 M_\odot$. The JHK magnitudes for these isochrones were transformed from the CIT system (Elias et al. 1982, 1983) to the 2MASS system using the transformations determined by Carpenter (2001). The relation for giant stars with $\log g < 2.0$ is taken from the 1.0 Gyr, solar metallicity Padova isochrone (Marigo et al. 2008; Bonatto, Bica & Girardi 2004)⁵. While the majority of stars lie in the expected dwarf range, a significant number of stars fall along the giant branch. Some of these stars may be rare carbon dwarfs, but the majority are most likely giants with inaccurate proper motion measurements in the PPMX catalog. Of the 2445 selected stars with $J - H > 0.8$ that have HATNet light curves, 87% have undetected proper motions or proper motions less than 10 mas/yr in the USNO-B1.0 catalog, this is compared to 28% of the sample with $J - H < 0.8$. A visual inspection of the POSS-I and POSS-II Digitized Sky Survey images for a number of the sources with $J - H > 0.8$ and $\mu > 100$ mas/yr revealed none with visually detectable proper motion, and in many cases the object consists of two close, comparably bright stars, for which misidentification of sources may be to blame for the spurious proper motion detection. This includes several stars where the PPMX and USNO-B1.0 proper motion values are comparable. We therefore apply an additional cut in the $J - H$ vs $H - K_S$ color-color diagram as shown in figure 1 to reduce the sample to 28,785 stars.

Figure 2 shows a $V - J$ vs. H_J reduced proper-motion (RPM; Luyten 1922) diagram for the sample. Here the RPM, H_J , is calculated as

$$H_J = J + 5 \log_{10}(\mu/1000) \quad (3)$$

and gives a rough measure of the absolute magnitude M_J of a star. We show roughly the lines separating main sequence dwarfs from sub-dwarfs and giants. In figure 3 we compare the RPM to M_J for 239 stars in the sample which have a Hipparcos parallax (Perryman et al. 1997). To remove additional giants from the sample we reject 1225 stars with $H_J < 3.0$, leaving our final sample of 27,560 stars.

3. SELECTION OF VARIABLE STARS

We use a number of techniques to search for light curves that show significant variability. The techniques include the phase-binning and harmonic-fitting Analysis of Variance periodograms (AoV/AoVHarm; Schwarzenberg-Czerny 1989, 1996), the Discrete Auto-Correlation Function (DACF; Edelson & Krolik 1988), and the Box-Least-Squares




FIG. 1.— $J - H$ vs. $H - K_S$ color-color diagram for 32,831 stars that have $V - K_S > 3.0$ with V taken either from the PPMX catalog (Tycho-2) or extrapolated from the 2MASS J , H and K_S magnitudes using eq. 1, $\mu > 30$ mas/yr from the PPMX catalog, and that have a HATNet light curve containing more than 1000 points (gray-scale points). The solid line shows the expected relation for cool dwarfs (Baraffe et al. 1998; Chabrier et al. 2000), while the dot-dashed line shows the expected relation for giants (Marigo et al. 2008; Bonatto, Bica & Girardi 2004). Stars outside the area enclosed by the dotted line are rejected. For display purposes we have added slight Gaussian noise to the observed colors in the plot.




FIG. 2.— $V - J$ vs. H_J RPM diagram for stars with $\mu > 30$ mas/yr passing $V - K_S$ and JHK color cuts. Note that in this plot we use V magnitudes that are transformed from the USNO-B1.0 photographic magnitudes for the majority stars, and not the V magnitudes transformed from JHK that were used in applying the initial $V - K_S$ cut. The lines separate main sequence dwarfs from sub-dwarfs and giants. We reject the 1225 stars with $H_J < 3.0$. The lines separating main sequence dwarfs from sub-dwarfs are taken from Yong & Lambert (2003).

(BLS; Kovács, Zucker & Mazeh 2002) algorithms as implemented in the VARTOOLS program⁶ (Hartman et al. 2008). We also conduct a search for flare-like events in the light curves. We apply the algorithms to both the EPD and TFA light curves of sources, when available. For the flare-search we use only the EPD light curves because the σ -clipping applied to the TFA light curves may remove real flares.

Because the light curves contain non-Gaussian, temporally-correlated noise, formal estimates for the

⁵ The isochrone was obtained from the CMD 2.1 web interface <http://stev.oapd.inaf.it/cgi-bin/cmd.2.1>

⁶ The VARTOOLS program is available at <http://www.cfa.harvard.edu/~jhartman/vartools/>

TABLE 1
COEFFICIENTS FOR TRANSFORMATIONS FROM USNO-B1.0 B_U AND R_U MAGNITUDES TO V (EQ. 2).

B_U	R_U	a	b	c	d	e	f	RMS [mag]
1	1	-0.76 ± 0.19	-0.02 ± 0.01	1.71 ± 0.15	-0.14 ± 0.01	0.15 ± 0.02	1.63 ± 0.52	0.27
1	2	0.72 ± 0.08	-0.061 ± 0.002	0.38 ± 0.06	-0.057 ± 0.006	0.114 ± 0.007	-0.89 ± 0.26	0.19
2	1	0.85 ± 0.09	-0.045 ± 0.006	0.44 ± 0.08	-0.051 ± 0.003	0.079 ± 0.008	-1.52 ± 0.24	0.22
2	2	0.73 ± 0.14	-0.164 ± 0.008	0.46 ± 0.13	-0.20 ± 0.01	0.35 ± 0.02	-0.75 ± 0.26	0.19




FIG. 3.— The absolute magnitude M_J vs. the RPM for 239 stars in the sample which have a Hipparcos parallax. This confirms that the RPM provides a rough estimate of the absolute magnitude for this sample of stars. We reject stars with $H_J < 3.0$, as these appear to be predominately giants.

variability detection significance are unreliable. We therefore have conducted Monte Carlo simulations of light curves with realistic noise properties to inform our choice of selection thresholds for several of the algorithms mentioned above, these are described in appendix A. In the following subsections we discuss the use of each of the variability selection algorithms in turn. We finish with a comparison of the different methods. The resulting catalog of variable stars is presented in appendix B.

3.1. AoV

The AoV periodogram is a method for detecting continuous periodic variations suggested by Schwarzenberg-Czerny (1989) which typically yields higher S/N detections than other periodograms. The original method suggested by Schwarzenberg-Czerny (1989) uses phase-binning for the model signal, so it is most comparable to the popular Phase Dispersion Minimization technique (PDM; Stellingwerf 1978). Following this Schwarzenberg-Czerny (1996) introduced an efficient method for fitting a Fourier series to a non-uniformly sampled light curve using a basis of orthogonal polynomials. When combined with an ANalysis Of VAriance (ANOVA) statistic, the result is the AoVHarm periodogram. We applied both methods to our light curves, and discuss each in turn.

3.1.1. Phase Binning AoV - Search for General Periodic Variability

We run the AoV algorithm with phase-binning on the full sample of stars. We first apply a 5σ iterative clipping to the light curve before searching for periods between 0.1 and 100 days. We use 8 phase bins, and generate the periodogram at a frequency resolution of $0.1/T$ where

T is the total time-span covered by a given light curve. We then determine the peak at 10 times higher resolution. As our figure of merit we use the signal-to-noise ratio (S/N), with a 5σ iterative clipping applied to the periodogram in calculating the noise (the RMS of the periodogram). Figure 4 shows the AoV S/N as a function of the peak period for various light curve subsamples. For comparison we also show the Monte Carlo simulation results for each subset. We adopt a separate selection threshold on S/N for each subsample. The thresholds have the form

$$S/N_{\min} = \begin{cases} S/N_0 & \text{if } P < P_0 \\ S/N_0 (P/P_0)^\alpha & \text{if } P \geq P_0 \end{cases} \quad (4)$$

The adopted values of S/N_0 , P_0 and α are listed for each subsample in Table 2. In addition to this selection we also reject detections with periods near 1 sidereal day, one of its harmonics, or other periods which appear as spikes in the histogram of detected periods (the latter includes periods between 5.71 and 5.80 days). For composite light curves that contain both 2K and 4K observations we take

$$S/N_{\min} = f_{2K} S/N_{\min,2K} + f_{4K} S/N_{\min,4K} \quad (5)$$

where f_{2K} and f_{4K} are the fraction of points in the light curve that come from 2K and 4K observations, and $S/N_{\min,2K}$ and $S/N_{\min,4K}$ are the 2K and 4K thresholds at the period of the composite light curve.

As summarized in table 3, our selection threshold passes a total of 1320 EPD light curves, and 1729 TFA light curves. These are inspected by eye to reject obvious false alarms and to identify EBs. There are 753 EPD light curves that we judge to show clear, continuous, periodic variability, 47 that show eclipses, 419 that we consider to be questionable (these are included in the catalog, but flagged as questionable), and 101 that we reject. For the TFA light curves the respective numbers are 1210, 64, 400, and 55. Note that the distinction between “clear” variability and “questionable” cases is fairly subjective. Generally we require the variations to be obvious to the eye for periods $\gtrsim 30$ days, for shorter periods we consider the selection to be questionable if it appears that the variability selection may be due to enhanced scatter on a few nights.

3.1.2. Harmonic AoV - Search for Sinusoidal Periodic Variability

The AoVHarm periodogram is generated for each star in a similar manner to the AoV periodogram. We run the algorithm using a sinusoid model with no higher harmonics (it is thus comparable to DFT methods, or to the popular Lomb-Scargle technique; Lomb 1976; Scargle 1982). As for the phase-binning AoV search we use eq. 5 for the selection threshold, with parameters given for each subsample in table 2. Figure 5 shows the AoVHarm S/N as

TABLE 2
SELECTION THRESHOLD PARAMETERS FOR AoV AND AoVHARM

Chip	LC Type	Mag.	S/N_0	AoV		α	AoVHarm		α
				P_0 [days]			S/N_0	P_0 [days]	
EPD	2K	6.5	30.0	5	0.40		40.0	4	0.41
EPD	2K	7.5	30.0	5	0.40		40.0	4	0.41
EPD	2K	8.5	30.0	5	0.40		40.0	4	0.41
EPD	2K	9.5	30.0	5	0.33		30.0	4	0.37
EPD	2K	10.5	25.0	5	0.23		30.0	4	0.37
EPD	2K	11.5	20.0	5	0.23		25.0	4	0.43
EPD	2K	12.5	20.0	5	0.23		25.0	4	0.36
EPD	2K	13.5	20.0	5	0.14		25.0	4	0.22
TFA	2K	6.5	20.0	5	0.14		30.0	4	0.16
TFA	2K	7.5	20.0	5	0.14		30.0	4	0.16
TFA	2K	8.5	20.0	5	0.14		30.0	4	0.16
TFA	2K	9.5	20.0	5	0.00		20.0	4	0.22
TFA	2K	10.5	15.0	5	0.00		20.0	4	0.13
TFA	2K	11.5	15.0	5	0.00		20.0	4	0.00
TFA	2K	12.5	15.0	5	0.00		20.0	4	0.00
TFA	2K	13.5	15.0	5	0.00		20.0	4	0.00
EPD	4K	7.5	30.0	5	0.33		30.0	4	0.50
EPD	4K	8.5	30.0	5	0.33		30.0	4	0.50
EPD	4K	9.5	30.0	5	0.33		30.0	4	0.50
EPD	4K	10.5	30.0	5	0.33		30.0	4	0.50
EPD	4K	11.5	25.0	5	0.23		25.0	4	0.43
EPD	4K	12.5	20.0	5	0.23		25.0	4	0.36
EPD	4K	12.5	20.0	5	0.23		20.0	4	0.43
EPD	4K	13.5	20.0	5	0.14		20.0	4	0.28
EPD	4K	14.5	20.0	5	0.00		20.0	4	0.22
TFA	4K	7.5	20.0	5	0.00		25.0	4	0.15
TFA	4K	8.5	20.0	5	0.00		25.0	4	0.15
TFA	4K	9.5	20.0	5	0.00		25.0	4	0.15
TFA	4K	10.5	20.0	5	0.00		25.0	4	0.06
TFA	4K	11.5	15.0	5	0.00		20.0	4	0.00
TFA	4K	12.5	15.0	5	0.00		20.0	4	0.00
TFA	4K	12.5	15.0	5	0.00		20.0	4	0.00
TFA	4K	13.5	15.0	5	0.00		20.0	4	0.00
TFA	4K	14.5	15.0	5	0.00		20.0	4	0.00

a function of peak period for several light curve subsamples.

Our selection threshold passes a total of 1337 EPD light curves and 1717 TFA light curves. Inspecting these by eye we find 1082 EPD light curves that show clear, continuous, periodic variability, 34 that show eclipses, 185 that we consider to be questionable, and 36 that we reject. For the TFA light curves the numbers are 1443, 43, 217, and 14 respectively (see also table 3).

3.2. DACF - Search for Quasi-Periodic Variability

The DACF is a technique that has frequently been used to determine the variation time-scale of quasi-periodic signals. In particular this technique is commonly applied in measuring the rotation period of a star from a light curve that exhibits variations due to the rotational modulation of starspots which may be varying in size, intensity, position, or number over time (e.g. Aigrain et al. 2008). It is not obvious, however, that the DACF method provides better period determinations than Fourier methods such as AoVHarm even in quasi-periodic cases. Since even in cases where the coherence time-scale is short compared to the period, the Fourier power spectrum should still have a predominant peak near the period of the star that can be determined in a straightforward fashion, whereas the automatic identification of the predominant variation time-scale from an autocorrelation function is non-trivial, as seen below. It is, therefore, with some skepticism that we attempt to use the DACF method to identify periods.

For each light curve we calculate the DACF at time lags ranging from 0 to 100 days with a step-size of 1 day. The binning time-scale of 1 day means that we will only be sensitive to variation time-scales $\gtrsim 2$ days. In practice, our peak finding algorithm limits us to periods $\gtrsim 10$ days. A light curve with a significant periodic or quasi-periodic signal with a timescale T will have a DACF that peaks at T , as well as at $2T$, $3T$, ... depending on the coherence of the signal.

To automate the selection of peaks in the DACF we use the following routine (below, y_i is the DACF value for time-lag t_i):

1. Identify connected sets of points (t_i, y_i) with $y_i > 0$.
2. Extend the left and right boundaries of each set until a local minimum is found in both directions.
3. Reject sets with 3 or fewer points, with 0 for the left boundary, or with the maximum time-lag computed for the right boundary. This leaves N_{peak} sets of points (peaks) to consider.
4. Fit a quadratic function $y(t)$ to each of the N_{peak} sets.
5. Letting χ^2_{N-3} and χ^2_{N-1} be the χ^2 values from fitting a quadratic function and a constant function respectively to the set, we perform an F-test on the statistic

$$f = \frac{(\chi^2_{N-1} - \chi^2_{N-3})/2}{\chi^2_{N-3}/(N-3)} \quad (6)$$

TABLE 3
SUMMARY OF VARIABLE STAR SELECTION

Variability Selection Type	LC Reduction Type	Automatically Selected	Non-EB Var.	EB	Quest.	Rej.
AoV	EPD	1320	753	47	419	101
AoV	TFA	1729	1210	64	400	55
AoV	Both ^a	781	576	41	53	5
AoV	Either ^b	2268	1387	70	766	151
AoV only ^c	Either ^b	450	120	4	317	79
AovHarm	EPD	1337	1082	34	185	36
AovHarm	TFA	1717	1443	43	217	14
AovHarm	Both ^a	936	806	24	27	5
AovHarm	Either ^b	2118	1719	53	375	45
AovHarm only ^c	Either ^b	450	120	4	317	79
BLS	EPD	463	79	60	39	281
BLS	TFA	444	160	81	39	160
BLS	Both ^a	155	62	52	4	13
BLS	Either ^b	752	177	89	74	428
BLS only ^c	Either ^b	439	1 ^d	21	64 ^d	428
DACF	EPD	1491	620	0	534	243
DACF	TFA	1190	507	0	318	95
DACF	Both ^a	353	203	0	27	21
DACF	Either ^b	2328	924	0	825	317
DACF only ^c	Either ^b	1344	373	0	582	528
Totals ^e		4579	2239	95	1176	1105

^a Stars that are selected by this method for both the EPD and TFA light curves. Note that stars that are automatically selected by this method for both EPD and TFA light curves but for which the by eye classification differs between the two light curve types will be included in the number of automatically selected variables, but will not be included in any of the subsequent columns for this row. ^b Stars that are selected by this method for either the EPD or the TFA light curves. ^c Stars that are selected exclusively by this method for either the EPD or the TFA light curves. For the Non-EB Var. and EB types stars are included in this row if they were only classified as that type during the visual inspection for this method. The questionable column lists the total number of stars that were flagged as questionable by this method and were either rejected during the visual inspection, or not selected, by all other methods. The rejected column lists the total number of stars that were rejected during the visual inspection for this method and did not pass the automatic selection for any of the other methods. ^d These stars are not included in the catalog of variables. ^e For the EB and non-EB variable classes we list the total number of stars that were classified as this type during the visual inspection for at least one method. Note that 36 stars are flagged as EBs during the visual inspection for one method and as non-EB variables during the visual inspection for another method, so a total of 2298 stars are flagged as either an EB or as a robust non-EB variable for at least one method. This total does not include flare stars selected in § 6.3, unless they are also identified as a variable by AoV, AovHarm, BLS or DACF. Including flare stars, the total number of stars with a robust variability detection is 2321. Of these, 1928 are not flagged as a probable blend or as having a problematic amplitude. For the questionable detections we list the total number of stars that are classified as questionable for at least one method and are not classified as a robust detection for any of the methods. For the rejections we list the total number of stars that were rejected during the visual inspection for all methods by which they were automatically selected.

to determine the significance of the quadratic fit relative to the constant function fit (see Lupton 1993). For each of the N_{peak} sets we record the time-lag of the peak and its error from the quadratic fit (t_p and σt_p), the peak DACF value and its error (y_p and σy_p), and the false alarm probability of the fit from the F-test (Pr_p). Note that this is not the false alarm probability of finding any connected set of points in the DACF that is well-fit by a quadratic in a random signal. In general, that false alarm probability will be higher.

- Starting from the peak with the shortest time lag t_p , identify the first peak with $Pr_p < Pr_{\text{lim}1}$ and $y_p > y_{\text{lim}}$ or with $Pr_p < Pr_{\text{lim}2}$, choose this peak as the period for the star. If there is no such peak in the DACF, then the star is not selected as a variable by this method. We adopt $Pr_{\text{lim}1} = 10^{-4}$; $Pr_{\text{lim}2}$ and y_{lim} are determined independently for

each subsample from the simulations. For a given subsample we take $Pr_{\text{lim}2}$ and y_{lim} to be the fifth smallest and largest values respectively from the simulations (i.e. the 99.5 percentile values).

In Figure 6 we plot the minimum Pr_p found in each DACF against the maximum y_p found and show the adopted cutoffs. We plot the results from the observed light curves and the simulations for four representative subsamples. Figure 7 shows an example of the DACF and selected peak for a periodic variable, and for one of the least significant detections that pass our selection. The un-phased light curves for each of these cases are also presented.

We find significant peaks in the DACF for a total of 1491 EPD light curves and 1190 TFA light curves. We inspect these by eye to eliminate obvious false alarms (typically cases where a light curve shows significant scatter on a few nights, often of several magnitudes or more),




FIG. 4.— Period vs. S/N from AoV for 4 representative light curve subsamples. On the left column we plot the observed values showing the light curves that pass the selection (dark filled points) and the light curves that do not pass the selection (grey filled points) separately, on the right column we plot the results from the Monte Carlo simulation for the corresponding subsample. The lines show the adopted S/N cut-off as a function of period. Note that in addition to the cut-off shown with the line, we also reject light curves for which the peak period is close to one sidereal day or a harmonic of one sidereal day.

we also note whether or not the detection appears to be robust and whether or not the period determination is likely to be accurate. For the EPD light curves we consider 465 of the detections to be robust and with a correct period, 155 to be robust but at the wrong period (typically the detected period is an integer multiple of the likely true period, see for example figure 7), 534 to be non-robust, and we reject 243 of the detections as clear false alarms. For the TFA light curves the respective numbers are 353, 154, 318, and 95 (see also table 3). The distinction between a robust and a non-robust detection is subjective, typically we consider a detection to be robust if the variability in the phased or unphased light curve is obvious to the eye and/or the DACF shows a set of clear regularly spaced peaks. There are some cases where the DACF shows clear regularly spaced peaks; however the scatter in the light curve appears to correlate with the phase. We consider many of these cases to be non-robust (most are light curves from 4K fields with periods near the lunar cycle). We include




FIG. 5.— Same as Figure 4, here we show the results for the AoVHarm period search.

all non-rejected detections in the final catalog together with flags indicating the reliability of the detection and the reliability of the period.

3.3. BLS - Search for Eclipses

The BLS algorithm, primarily used in searches for transiting planets, detects periodic box-like dips in a light curve. This algorithm may be more sensitive to detached binaries with sharp-featured light curves than the other methods used. For our implementation of the BLS algorithm we search 10,000 frequency points within a period range of 0.1 to 20.0 days. To search for long period events where the eclipse duration may be only a very short fraction of the orbital period, we repeated the search at a higher frequency resolution using 100,000 points within a period range of 1.0 to 20.0 days. At each trial frequency we bin the phased light curve into 200 bins, and search over fractional eclipse durations ranging from 0.01 to 0.1 in phase. Figure 8 shows the S/N vs the period for the EPD and TFA light curves. We select 752 stars with $S/N > 10.0$ and with a period not close to 1 sidereal day or a harmonic of a sidereal day as potentially eclipsing systems. We do not use the Monte Carlo simulations to set the selection thresholds for this method because distinguishing between red noise and an eclipse signal by eye is less ambiguous than distinguishing between red




FIG. 6.— Minimum false alarm probability Pr_p vs. maximum peak value y_p found in each DACF for 4 representative light curve subsamples. On the left column we plot the observed values showing the light curves that pass the selection (dark filled points) and the light curves that do not pass the selection (grey filled points) separately, on the right column we plot the results from the Monte Carlo simulation for the corresponding subsample. The lines show the adopted Pr_p/y_p cut-off. Note that the plotted Pr_p and y_p values for a given DACF may not come from the same peak. In selecting peaks from the DACF we require that both the Pr_p and the y_p values for a given peak pass the selection. This plot is meant only to provide an approximate visualization of the selection.

noise and general periodic or quasi-periodic variability. The selected light curves are inspected by eye to identify eclipsing systems. A total of 89 candidate EB systems are found in this manner, 8 are found in the EPD light curves only, 29 are found in the TFA light curves only, and 52 are found in both the EPD and TFA light curves (see also table 3).

3.4. Search for Flares

As noted in the introduction, flaring is a common phenomenon among K and M dwarfs. While we were inspecting the light curves of candidate variable stars we noticed a number of stars showing significant flares. We therefore decided to conduct a systematic search for flare events in the light curves. Most optical stellar flares show a very steep rise typically lasting from a few seconds to several minutes. Krautter (1996) notes that flares can be divided into two classes based on their decay times: “impulsive” flares have decay times of a few minutes, to a

few tens of minutes, while “long-decay” flares have decay times of up to a few hours. Due to the 5-minute sampling of the HATNet light curves, flares of the former type will only affect one or two observations in a light curve, while flares of the latter type might affect tens of observations. In general it is very difficult to determine whether a given outlier in a light curve is due to a flare or bad photometry without inspecting the images from which an observation was obtained. This is impractical to do for tens of thousands of light curves when each light curve may contain tens to hundreds of outliers. While observations that are potentially corrupted are flagged, in practice the automated routines that generate these flags do not catch all cases of bad photometry. We therefore do not attempt to identify individual “impulsive” flares in the light curves, and instead conduct a statistical study of the frequency of these flares (§ 6.3). Long-decay flares, on the other hand, may be searched for in an automated fashion if a functional form for the decay is assumed (this is similar to searching for microlensing events, see for example Nataf et al. 2009). To search for long-decay flares we used the following algorithm:

1. Compute m_0 , the median magnitude of the light curve, and δ_0 the median deviation from the median.
2. Identify all sets of consecutive points with $m - m_0 < -3\delta_0$. Let t_0 be the time of the brightest observation in a given set, and let N be the number of consecutive points following and including t_0 with $m - m_0 < -2\delta_0$. We proceed with the set if $N > 3$.
3. Use the Levenberg-Marquardt algorithm (Marquardt 1963) to fit to the N points a function of the form:

$$m(t) = -2.5 \log_{10} \left(A e^{-(t-t_0)/\tau} + 1 \right) + m_1 \quad (7)$$

where A , τ and m_1 are the free parameters. Here A is the peak intensity of the flare relative to the non-flaring intensity, τ is the decay timescale, and m_1 is the magnitude of the star before the flare. For the initial values we take $m_1 = m_0$, $\tau = 0.02$ days, and $A = 10^{-0.4(m_p - m_0)} - 1$, where m_p is the magnitude at the peak.

4. Perform an F-test on the statistic given in eq. 6 where χ^2_{N-3} in this case is the χ^2 value from fitting eq. 7. If the false alarm probability is greater than 1%, reject the candidate. If not, increase the number of points by one and repeat step 3. Continue as long as the false alarm probability decreases.
5. We reject any flare candidate for which there are at least two other candidate flares from light curves in the same field that occur within 0.1 days of the flare candidate.
6. Let the number of points with $t < t_0$ and $t_0 - t < 0.05$ days be N_{before} and the number of points with $t > t_1$ and $t - t_1 < 0.05$ days be N_{after} . Here t_1 is the time of the last observation included in the fit. Reject the candidate flare if $N_{\text{before}} < 2$ or $N_{\text{after}} < 2$. Also reject the candidate flare if $A < 0.$,




FIG. 7.— Top: Examples of the DACF for one of the highest significance detections of variability (left) and for one of the lowest significance detections (right). In each case the dark line shows the quadratic fit to the DACF used to determine the period of variation. Bottom: Un-phased light curves for the two stars. Note that the period for the star on the right is likely half the value determined by the peak identification algorithm. In this case the peak at $P \sim 10$ days did not pass the cut on Pr_p , while the second peak at $P \sim 20$ days did. The $P \sim 10$ days signal is the top peak in the AoV and AoVHarm periodograms for this star, however the S/N for both periodograms is below our selection threshold.

$A > 10.0$, $\tau < 0.001$ days, $\tau > 0.5$ days, $A < \sigma_A$, $\tau < \sigma_\tau$, or if the false alarm probability from step 4 is greater than 0.1%. Here σ_A and σ_τ are the formal uncertainties on A and τ respectively. The selection on A is used to reject numerous light curves with significant outliers which appear to be due to artifacts in the data rather than flares.

We apply the above algorithm to the non-composite EPD light curves (i.e. light curves from each field are processed independently for stars with light curves from multiple fields). The algorithm is applied both on the raw EPD light curves, and on EPD light curves that are high-pass filtered by subtracting from each point the median of all points that are within 0.1 day of that point. There are a total of 23,589 stars with EPD light curves that are analyzed, we exclude from the analysis 4 stars from the full sample for which only σ -clipped TFA light curves are available. A total of 320 candidate flare events from 281 stars are selected. These are inspected by eye to yield the final sample of 64 flare events from 60 stars. Figure 9 shows two examples of these large-amplitude, long-decay flares. The identified flares have peak intensities that range from $A = 0.09$ to $A = 4.21$ and decay time-scales that range from $\tau = 4$ minutes to $\tau = 1.7$ hours.

3.5. Comparison of Selection Methods

As seen in Table 3 more stars pass the automatic selections for AoV or for DACF than pass the automatic selections for AoVHarm. On the other hand, there are

more robust detections found by AoVHarm than by the other methods. The latter result may be due, in part, to a bias toward sinusoidal signals in the by-eye selection. However, taking it at face value, it appears that the AoVHarm method is a more robust period-finder for this sample of light curves than either the AoV or the DACF methods. For the 1075 TFA light curves that are classified as robust detections during the visual inspections for both the AoV and AoVHarm methods, the S/N for the AoVHarm detection is greater than the S/N for the AoV detection in all but 12 cases. This is in line with the long-known fact that Fourier-based period-finding methods generally yield higher S/N detections than phase-binning methods, even in cases where the signal is non-sinusoidal (Kovács 1980).

A direct comparison between the DACF and the AoVHarm or AoV methods is difficult since the selection for DACF is quite different from the selections for AoV or AoVHarm. However, it is apparent from table 3 that, at least for our selection thresholds, the DACF detections are generally found to be less reliable than the AoVHarm or AoV detections. Again there may be a bias in the by-eye selection against the types of variables that pass DACF, however it is also likely that the increased complexity in identifying a period from the auto-correlation function results in more false alarms than are generated by the periodogram-based methods.

As expected, BLS appears to be the best technique for identifying eclipsing binaries; 89 out of the 95 potential




FIG. 8.— Period vs. S/N from BLS for EPD and TFA light curves. The dark points show light curves that pass the $S/N > 10$ cut and do not have a period near a sidereal day or one of its harmonics. The grey points show light curves that do not pass this cut.




FIG. 9.— Two examples of large-amplitude, long-decay flares seen in the HATNet light curves. In each case the solid line shows the fit of eq. 7 to the light curve that was done in automatically identifying candidate flares.

eclipsing binaries identified in our survey are selected by BLS, including 21 that are identified exclusively by BLS. The second most successful method for identifying EBs is the AoV method which identified 70 of the 95 potential eclipsing binaries, and exclusively identified 4 of them.

4. VARIABILITY BLENDING

While the wide FOV of the HATNet telescopes allows a significant number of bright stars to be simultaneously observed, the downside to this design is that the pixel scale is necessarily large, so a given light curve often in-

cludes flux contributions from many stars. Blending is a particularly significant issue in high stellar density fields near the Galactic plane. A star blended with a nearby variable star may be incorrectly identified as a variable based on its light curve. If the stars are separated by more than a pixel or two, it may be possible to distinguish the real variable from the blend by comparing the amplitudes of their light curves. However, because photometry is only obtained for stars down to a limiting magnitude (the value used varies from field to field), in many cases we do not have light curves for all the faint neighbors near a given candidate variable star, so we cannot easily determine which star is the true variable. In these cases we can still give an indication of whether or not a candidate is likely to be a blend by determining the expected flux contribution from all neighboring stars to the candidate's light curve.

To determine whether or not a candidate is blended with a nearby variable star that has a light curve, we measure the peak-to-peak light curve amplitude (in flux) of all stars within $2'$ of the candidate. If any neighbor has an amplitude that is greater than twice the flux amplitude of the candidate, the candidate is flagged as a probable blend. If any neighbor has an amplitude that is between half and twice the flux amplitude of the candidate, the candidate is flagged as a potential blend. If any neighbor has an amplitude that is between 10% and half the flux amplitude of the candidate, the candidate is flagged as an unlikely blend. And finally we flag the candidate as a non-blend if all neighbors have amplitudes that are less than 10% that of the candidate. We determine the amplitude of a light curve by fitting to it 10 different Fourier series of the form:

$$m(t) = m_0 + \sum_{i=1}^N a_i \sin(2\pi i t/P + \phi_i) \quad (8)$$

with N ranging from 1 to 10. Here P is the period of the light curve. We perform an F-test to determine the significance of each fit relative to fitting a constant function to the light curve, and choose the amplitude of the Fourier series with the lowest false alarm probability. If the lowest false alarm probability is greater than 10% we set the amplitude to zero. We try all periods identified for each candidate by the variability searches described in § 3, and adopt the largest amplitude found. For candidates that have light curves from multiple fields, or that have both ISM and AP reductions, we do the amplitude test on each separate field/reduction and adopt the most significant blending flag found for the candidate. If the amplitude of the candidate variable star is set to zero for a given field/reduction we do not use that field/reduction in determining the blending flag. If this is true for all fields/reductions we flag the candidate as test.

To determine whether or not a candidate is potentially blended with a nearby faint variable star that does not have a light curve, we compare the observed amplitude of the candidate to its expected amplitude if a neighboring star were variable with an intrinsic amplitude of 1.0 mag. We assume that an amplitude of 1.0 mag is roughly the maximum value that one might expect for a short period variable star. If the measured amplitude is less than the

expected amplitude then we flag the candidate as a potential blend, if it is greater than the expected amplitude and less than twice the expected amplitude we flag the candidate as an unlikely blend, and if it is greater than twice the expected amplitude, we flag it as a non-blend. The test is done for all stars within $2'$ of the candidate that do not have a light curve, and we adopt the most significant blending flag found for the candidate. To determine the expected amplitude of the candidate star induced by the neighbor, we note that a star with magnitude m_1 located near a variable star with magnitude m_2 and amplitude $\Delta m_2 > 0$, has an expected light curve amplitude that is given by

$$\Delta m_{1,AP} = 2.5 \log_{10} \left[f_1 10^{-0.4m_1} + f_2 10^{-0.4(m_2 - \Delta m_2)} \right] - 2.5 \log_{10} \left[f_1 10^{-0.4m_1} + f_2 10^{-0.4m_2} \right] \quad (9)$$

for the case of aperture photometry, and by

$$\Delta m_{1,ISM} = 2.5 \log_{10} \left[f_1 10^{-0.4m_1} + f_2 10^{-0.4m_2} (10^{0.4\Delta m_2} - 1) \right] - 2.5 \log_{10} \left[f_1 10^{-0.4m_1} \right] \quad (10)$$

for the case of image subtraction photometry. The two expressions differ because in the ISM pipeline photometry is done on difference images (only differential flux is summed in the aperture), whereas in the AP pipeline photometry is done directly on the science images (all stellar flux is summed in the aperture). Here $f_{1,2}$ is the fraction of the flux from star 1(2) that falls within the aperture, and we use the catalog values (transformed from 2MASS) for m_1 and m_2 . To determine f_1 and f_2 we integrate the intersection between the circular aperture and a Gaussian PSF which we assume to have a FWHM of 2 pixels (this is a typical effective “seeing” for both the 2K and 4K images), we do not consider pixelation effects in making this estimate.

We compare the results from the two blending tests for each candidate, and adopt the most significant blending flag from among the two tests for the catalog. We do not run the test on the candidate flare stars, unless the star was selected as a variable by another method as well. Out of the 3474 stars that are in either the first or the second catalog, 936 are flagged as unblended, 451 are flagged as unlikely blends, 1397 are flagged as potential blends, 399 are flagged as probable blends, and 291 are found to have problematic amplitudes (cases where the amplitude measuring algorithm failed for the star in question).

5. MATCH TO OTHER CATALOGS

5.1. Match to Other Variable Star Surveys

We match all 3496 stars selected as potential variables to the combined General Catalogue of Variable Stars (GCVS; Samus’ et al. 2006), the New Catalogue of Suspected Variable Stars (NSV; Kholopov 1982) and its supplement (NSVS; Kazarovets, Samus’ & Durlevich 1998)⁷. We also match to the ROTSE catalog of variable stars (Akerlof et al. 2000), to the ALL Sky Automated Survey Catalogue of Variable Stars (ACVS;

Pojmanski 2002)⁸, and to the Super-WASP catalogue of periodic variables coincident with ROSAT X-ray sources (Norton et al. 2007). In all cases we use a $2'$ matching radius. We use a large matching radius to include matches to known variables that may be blended with stars in our sample. In total 77 of our candidate variables lie within $2'$ of a source in one of these catalogs, meaning that 3419 are new identifications. This includes 36 that match to a source in the GCVS, 4 that match to a source in the NSV, 7 that match to a source in the NSVS, 4 that match to a source in the ACVS, 8 that match to a source in the ROTSE catalog (4 of which are in the GCVS as well), and 23 that match to a Super-WASP source (2 of these are in their catalogue of previously identified variables). Two of the 36 candidate variables that match to a source in the GCVS (HAT-215-0001451 and HAT-215-0001491) actually match to the same source, V1097 Tau, a weak emission-line T Tauri star. Both stars are flagged as probable blends in our catalog, in this case HAT-215-0001491 is the correct variable while HAT-215-0001451 is the blend.

We inspect each of the 77 candidates with a potential match and find that the match is incorrect for 39 of them and correct for 38. For 23 of the 39 incorrect matches the candidate variable is flagged as a probable blend in our catalog. In 8 cases the candidate variable is flagged as a potential blend, in 4 cases it is flagged as an unlikely blend, in 3 cases it is flagged as unblended, and in 1 case the amplitude is considered problematic. The match appears to be correct for four of the candidate variables flagged as probable blends. In addition to HAT-215-0001491, the stars HAT-239-0000221 and HAT-239-0000513 both match correctly to sources in the GCVS. These stars form a common proper motion, low mass binary system. Both stars are flagged as probable blends in our catalog. Each matches separately, and correctly, to a flare star in the GCVS (V0647 Her and V0639 Her respectively). For the NSV-matching probable blend candidate HAT-121-0003519 the match may be correct, though the positional uncertainty of the NSV source is high. A variable star classification is not available for this source.

The 16 variables that match correctly to a source in the GCVS include 4 BY Draconis-type rotational variables, 6 UV Ceti-type flare stars, 1 INT class Orion variable of the T Tauri type, and 5 eclipsing systems. The EBs include the two W UMa-type contact systems DY CVn and V1104 Her, the two Algol-type systems DK CVn and V1001 Cas, and the M3V/white dwarf EB DE CVn (van den Besselaar et al. 2007).

Table 16, at the end of the paper, lists the first ten cross-identifications, the full table is available electronically with the rest of the catalog.

5.2. Match to ROSAT

We match all 3496 stars selected as potential variables to the ROSAT All-sky survey source catalog (Voges et al. 1999) using the US National Virtual Observatory catalog matching utilities. We use a 3.5σ positional matching criterion. A total of 248 of the variables match to an X-ray source, including 237 stars in our primary catalog

⁷ The GCVS, NSV and NSVS were obtained from <http://www.sai.msu.su/groups/cluster/gcvs/gcvs/> on 2009 April 7

⁸ Version 1.1 obtained from <http://www.astrouw.edu.pl/asas/?page=catalogues>

of periodic variables, 14 of the EBs, and 24 of the 60 flare stars. A few of the variable stars are close neighbors where one is likely to be a blend of the other, so there are 243 distinct X-ray sources that are matched to. Table 17 at the end of the paper gives the cross-matches. The full table is available electronically with the rest of the catalog. In Section 6.2 we discuss the X-ray properties of the rotational variables.

6. DISCUSSION

6.1. Eclipsing Binaries

The 95 stars that we identify as potential EBs have periods ranging from $P = 0.193$ days to $P = 24.381$ days. We flag 11 of the candidate EBs as probable blends, 23 as potential blends, 21 as unlikely blends, 35 as unblended, and 5 as having problematic amplitudes. Figure 10 shows phased light curves for 12 of the EBs.




FIG. 10.— Example phased light curves for 12 of the 95 potential EBs found in the survey. The period listed is in days.

In addition to matching the candidate EBs to other variable star catalogs (§ 5.1) and to ROSAT (§ 5.2) we also checked for matches to previously studied objects using SIMBAD. The following objects had noteworthy matches:

1. *HAT-148-0000574*: matches to the X-ray source 1RXS J154727.5+450803, and was previously discovered to be an SB2 system by Mochnacki et al. (2002), we discuss this system in detail in § 6.1.1.
2. *HAT-216-0002918* and *HAT-216-0003316*: match to CCDM J04404+3127A and CCDM

J04404+3127B, respectively, which form a common proper-motion 15'' binary system. The two stars are both selected as candidate EBs with the same period, and are both flagged as potential blends in the catalog; based on a visual inspection of the light curves we conclude that the fainter component (HAT-216-0003316) is most likely the true $P = 2.048$ day EB. The fainter component, which has spectral type M3 on SIMBAD, also matches to the X-ray source RX J0440.3+3126.

3. *HAT-127-0008153*: matches to CCDM J03041+4203B, which is the fainter component in a common proper-motion 20'' binary system. This star is flagged as a potential blend, the brighter component appears to match to the X-ray source 1RXS J030403.8+420319. Based on a visual inspection of the light curves we conclude that HAT-127-0008153 is likely the true variable.
4. *HAT-169-0003847*: is 24'' from the Super-WASP variable 1SWASP J034433.95+395948.0, the two stars are blended in the HATNet images, however from a visual inspection of the light curves we conclude that HAT-169-0003847 is likely the true variable.
5. *HAT-192-0001841*: is 46'' from a high proper-motion, K0 star BD+41 2679. The two stars may be members of a common proper-motion binary system (the former has a proper motion of 65.79, -151.77 mas/yr in RA and DEC respectively, while the latter has 89.76, -117.14 mas/yr).
6. *HAT-169-0003847*: is flagged as an unlikely blend, and we confirm that BD+41 2679 is not the true variable.
7. *HAT-193-0008020*: matches to GSC 03063-02208 and has a spectral type of M0e listed on SIMBAD (see also Mason et al. 2000).
8. *HAT-216-0007033*: matches to the X-ray source RX J0436.1+2733 and has spectral type M4 listed on SIMBAD.
9. *HAT-341-0019185*: is 43'' from TYC 1097-291-1, the two stars appear to be members of a common proper motion binary system. HAT-341-0019185 is flagged as a probable blend in our catalog, though it does not appear that TYC 1097-291-1 is the real variable.

6.1.1. The Low-mass EB 1RXS J154727.5+450803

The EB HAT-148-0000574 matches to 1RXS J154727.5+450803. Using RV observations obtained with the Cassegrain spectrograph on the David Dunlap Observatory (DDO) 1.88 m telescope⁹, Mochnacki et al. (2002) found that this object is a $P = 3.54997 \pm 0.00005$ day double-lined spectroscopic binary system with component masses $\gtrsim 0.26 M_{\odot}$. This system, however, was not previously known to be eclipsing. Here we combine the published RV curves

⁹ Based on data obtained at the David Dunlap Observatory, University of Toronto

from Mochnecki et al. (2002) with the HATNet I-band light curve to provide preliminary estimates for the masses and radii of the component stars.

Figure 11 shows the EPD HATNet light curve phased at a period of $P = 3.550018$ days together with a model fit, while figure 12 shows a fit to the radial velocity observations taken from Mochnecki et al. (2002). Note the out of eclipse variations in the light curve, presumably due to spots on one or both of the components, which indicates that the rotation period of one or both of the stars is tidally locked to the orbital period. Since the HATNet light curve is not of high enough quality to measure the radii to better than a few percent precision, we do not fit a detailed spot model to the light curve, and instead simply fit a harmonic series to the out of eclipse observations and then subtract it from the full light curve. We model the light curve using the JKTEBOP program (Southworth, Maxted & Smalley 2004; Southworth et al. 2004) which is based on the Eclipsing Binary Orbit Program (EBOP; Popper & Etzel 1981; Etzel 1981; Nelson & Davis 1972), but includes more sophisticated minimization and error analysis routines. We used the DEBiL program (Devor 2005) to measure the eclipse minimum times from the light curve, which in turn were used with the RV curves to determine the ephemeris. In modeling the RV curves we fix $e = 0$ (Mochnecki et al. 2002, found $e = 0.008 \pm 0.007$), and we fix $k = R_2/R_1 = 1.0$ in modelling the light curve given $q = 1.00 \pm 0.02$ from the fit to the RV curves (the light curve is not precise enough to provide a meaningful constraint on k). For completeness we note that we assumed quadratic limb darkening coefficients of $a = 0.257$, $b = 0.586$ for both stars (Claret 2000), which are appropriate for a $T_{\text{eff}} = 3000$ K, $\log g = 4.5$, solar metallicity star. The results are insensitive to the adopted limb darkening coefficients; we also performed the fit using the coefficients appropriate for a $T_{\text{eff}} = 4000$ K, $\log g = 4.5$ star and found negligible differences in the resulting parameters and uncertainties. The parameters for the system are given in table 4. Note that the 1σ errors given on the masses and radii are determined from a Monte Carlo simulation (Southworth et al. 2005). These are likely to be overly optimistic given the inaccurate treatment of the spots, and our assumption that the component radii are equal.

Table 5 lists the masses and radii of the 4 other known double-lined detached EBs with at least one main sequence component that has a mass less than $0.3M_{\odot}$. We do not include RR Cae which is a white-dwarf/M-dwarf EB that has presumably undergone mass transfer (Maxted 2007). In figure 13 we plot the mass-radius relation for stars in the range $0.15M_{\odot} < M < 0.3M_{\odot}$. Like the components of CM Dra and the secondary of GJ 3236, and unlike the components of SDSS-MEB-1 and the secondary of 2MASSJ04463285+190432, the components of 1RXS J154727.5+450803 have radii that are larger than predicted from the Baraffe et al. (1998) isochrones (if the age is $\gtrsim 200$ Myr). The radii are $\sim 10\%$ larger than the predicted radius in the 1.0 Gyr, solar metallicity isochrone. High precision photometric and spectroscopic follow-up observations, and a more sophisticated analysis of the data are needed to confirm this.

6.2. Rotational Variables

TABLE 4
PARAMETERS FOR THE EB 1RXS J154727.5+450803

Coordinates and Photometry	
RA (J2000)	15:47:27.42 ^a
DEC (J2000)	+45:07:51.39 ^a
Proper Motions [mas/yr]	-259, 200 ^b
J	9.082 mag ^c
H	8.463 mag ^c
K	8.215 mag ^c
Ephemerides	
P	3.5500184 $\pm$ 0.0000018 day
HJD	2451232.89534 $\pm$ 0.00094
Physical Parameters	
M_1	0.2576 $\pm$ 0.0085 $M_{\odot}$
M_2	0.2585 $\pm$ 0.0080 $M_{\odot}$
$R_1 = R_2$	0.2895 $\pm$ 0.0068 $R_{\odot}$
RV Fit Parameters	
γ	-21.21 $\pm$ 0.41 km/s
K_1	55.98 $\pm$ 0.76 km/s
K_2	55.78 $\pm$ 0.83 km/s
e	0.0 ^d
LC Fit Parameters	
J_2/J_1	1.0734 $\pm$ 0.030
$(R_1 + R_2)/a$	0.0737 $\pm$ 0.0014
R_2/R_1	1.0 ^e
i	86.673 $^{\circ}$ $\pm$ 0.068 $^{\circ}$

^a SIMBAD^b PPMX^c 2MASS^d fixed^e fixed based on $q = 1.004 \pm 0.020$ from fitting the RV curves.

Figure 14 shows example phased light curves for 12 randomly selected rotational variables found in our survey. For the following analysis we only consider stars in our variables catalog that are identified as reliable detections for at least one search algorithm and that are not flagged as probable blends or as having problematic amplitudes. In order of preference, we adopt the AoVHarm TFA, AoVHarm EPD, AoV TFA, AoV EPD, DACF TFA or DACF EPD period for the star, choosing the period found by the first method for which the detection is considered secure.

6.2.1. Period-Amplitude Relation

For FGK stars there is a well-known anti-correlation between stellar activity measured from emission in the H and K line cores, or from H α emission, and the Rossby number (R_O , the ratio of the rotation period to the characteristic time scale of convection, see Noyes et al. 1984), which saturates for short periods. Similar anti-correlations with saturation have been seen between R_O and the X-ray to bolometric luminosity ratio (e.g. Pizzolato et al. 2003), and between R_O and the amplitude of photometric variability (e.g. Messina et al. 2001; Hartman et al. 2009). Main sequence stars with $M \lesssim 0.35M_{\odot}$ are fully convective (Chabrier & Baraffe 1997), so one might expect that the rotation-activity relation breaks down, or significantly changes, at this mass. Despite this expectation, several studies have indicated that the rotation-activity relation (measured using $v \sin i$ and H α respectively) continues for late M-dwarfs (Delfosse et al. 1998; Mohanty & Basri 2003; Reiners & Basri 2007). Recently, however, West & Basri

TABLE 5
OTHER DOUBLE-LINED EBs WITH A VERY LATE M DWARF COMPONENT

Name	Period [days]	$M_1 [M_\odot]$	$M_2 [M_\odot]$	$R_1 [R_\odot]$	$R_2 [R_\odot]$
CM Dra	1.27	0.2310 ± 0.0009	0.2141 ± 0.0010	0.2534 ± 0.0019	0.2396 ± 0.0015
SDSS-MEB-1	0.407	0.272 ± 0.020	0.240 ± 0.022	0.268 ± 0.010	0.248 ± 0.009
2MASSJ04463285+190432	0.619	0.47 ± 0.05	0.19 ± 0.02	0.57 ± 0.02	0.21 ± 0.01
GJ 3236	0.771	0.376 ± 0.016	0.281 ± 0.015	0.3795 ± 0.0084	0.300 ± 0.015

REFERENCES. — CM Dra: Morales et al. (2009); Lacy (1977); Metcalfe et al. (1996); SDSS-MEB-1: Blake et al. (2008); 2MASSJ04463285+190432: Hebb et al. (2006); GJ 3236: the parameters listed for this system are determined by giving equal weight to the three models in Irwin et al. (2009b)




FIG. 11.— Phased HATNet I-band light curve for the low-mass EB 1RXS J154727.5+450803. The top panel shows the full EPD light curve, the bottom two panels show a model fit to the two eclipses after subtracting a harmonic series fit to the out of eclipse portion of the light curve.

(2009) have found that rotation and activity may not always be linked for these stars.

In figure 15 we plot the rotation period against the peak-to-peak amplitude for stars in several color bins. The peak-to-peak amplitude is calculated for the EPD light curves as described in section 4; for stars observed in multiple fields we take the amplitude of the combined light curve. Stars without an available EPD light curve, or for which the amplitude measuring algorithm failed, are not included in the plot. A total of 1525 stars are included in the plot. For stars with $V - K_S < 5.0$ (corresponding roughly to $M \gtrsim 0.25 M_\odot$) the photometric amplitude and the period are anti-correlated at high significance. There appears to be a cut-off period, such that




FIG. 12.— Circular orbit fit to the RV curves for the low-mass EB 1RXS J154727.5+450803. The observations are taken from Mochnacki et al. (2002).




FIG. 13.— Mass-radius relation for 7 main sequence stars in double-lined DEBs with $M < 0.3 M_\odot$. The two points with the smallest error bars are the components of CM Dra, the open circles are the components of SDSS-MEB-1, the open square is the secondary component of 2MASSJ04463285+190432, the X is the secondary component of GJ 3236, and the filled square marks the two components of 1RXS J154727.5+450803. The solid line shows the 1.0 Gyr, solar metallicity isochrone from Baraffe et al. (1998). Note that the error bars for 1RXS J154727.5+450803 do not incorporate systematic errors that may result from not properly modelling the spots or allowing the stars to have unequal radii.

the period and amplitude are uncorrelated for stars with periods shorter than the cut-off, and are anti-correlated for stars with periods longer than the cut-off. In other words, the relations are saturated below a critical rotation period. Hartman et al. (2009) find a saturation threshold of $R_O = 0.31$, which for a $0.6 M_\odot$ star corresponds to a period of ~ 8 days (assuming $(B-V)_0 = 1.32$




FIG. 14.— Phased EPD light curves for 12 randomly selected rotational variables found in the survey. The period listed is in days. The zero-point phase is arbitrary. For this figure we use the period found by applying AoVHarm to the TFA light curves.

for stars of this mass, and using the empirical relation between the convective time scale and $(B - V)_0$ from Noyes et al. 1984). This is consistent with what we find for the bluest stars in our sample. For stars with $V - K_S > 5.0$ ($M \lesssim 0.25 M_\odot$), the period and amplitude are not significantly correlated, at least for periods $\lesssim 30$ days. This result suggests that the distribution of starspots on late M dwarfs is uncorrelated with rotation period over a large period range, and is perhaps at odds with $H\alpha/v \sin i$ studies which indicate a drop in activity for very late M-dwarf stars with $v \sin i \lesssim 10$ km/s (e.g. Mohanty & Basri 2003).

6.2.2. Period-Color/Period-Mass Relation

In figure 16 we compare the distribution of $V - K_S$ colors for periodic variables to the distribution for all stars in the sample. We also show the fraction of stars that are variable with peak-to-peak amplitudes greater than 0.01 mag as a function of $V - K_S$.

The plotted relation has been corrected for completeness by conducting sinusoid injection/recovery simulations to estimate our detection efficiency. In conducting these tests we divide the sample into 90 period/amplitude/color bins. We use color bins of $2.0 < V - K_S < 3.5$, $3.5 < V - K_S < 4.0$, $4.0 < V - K_S < 4.5$, $4.5 < V - K_S < 5.0$ and $5.0 < V - K_S < 6.0$, three period bins of 0.1-1 day, 1-10 days, and 10-100 days, and 5 amplitude bins logarithmically spanning 0.01 to 1.0 mag. While ideally a much finer grid would be used for these simulations, we were limited by computational




FIG. 15.— Rotation period vs. peak-to-peak photometric amplitude for 1525 stars. We divide the sample into 4 bins by $V - K_S$ color, corresponding roughly to $M \gtrsim 0.6 M_\odot$, $0.5 M_\odot \lesssim M \lesssim 0.6 M_\odot$, $0.25 M_\odot \lesssim M \lesssim 0.5 M_\odot$, and $M \lesssim 0.25 M_\odot$, from blue to red. We also list the Spearman rank-order correlation coefficient and the statistical significance of the correlation for each sample (note that negative values of r_s imply that the period and amplitude are anti-correlated). For stars with $V - K_S < 5.0$ the period and peak-to-peak amplitude are anti-correlated with high significance. The relation appears to be saturated for periods $\lesssim 5$ days, with hints that the saturation period increases for decreasing stellar mass. For stars with $V - K_S > 5.0$ ($M \lesssim 0.25 M_\odot$), the period and amplitude are not significantly correlated for $P \lesssim 30$ days.

resources to this fairly coarse sampling. For each bin we randomly select 1000 stars with the appropriate color (for color bins with fewer than 1000 stars we select with replacement). For each selected star we then choose a random period and amplitude drawn from uniform-log distributions over the bin and inject a sine curve with that period/amplitude and a random phase into the light curve of the star. If both EPD and TFA light curves are available for the star we inject the same signal into both light curves. We do not reduce the amplitude for the injection into the TFA light curve, this may cause us to slightly overestimate our detection efficiency. If the star was identified as a variable or a potential variable by our survey, we first remove the true variable signal from the light curve by fitting a harmonic series to the phased light curve before injecting the simulated signal. We then process the simulated signals through the AoV, AoVHarm and DACF algorithms using the same selection parameters as used for selecting the real variables. We do not apply by-eye selections on the simulated light curves, so our detection efficiency may be overestimated (particularly for longer period stars where the by-eye selection tended to be stricter than the automatic cuts). To get the completeness corrected variability fraction we weight each real detected variable by $1/f$ where f is the fraction of simulated signals that are recovered for the period/amplitude/color bin that the real variable falls in. We find that we are roughly $\sim 80\%$ complete over our sample of stars for peak-to-peak amplitudes greater than

0.01 mag and periods between 0.1 and 100 days. Considering the recovery fraction separately for each variability method, we find that $\sim 91\%$ of the simulations are recovered by AoVHarm, and in $\sim 97\%$ of the recovered simulations the recovered frequency is within 0.001 day^{-1} of the injected frequency. For AoV we again find that $\sim 91\%$ of the simulations are recovered, but that fraction of recovered simulations where the recovered frequency is within 0.001 day^{-1} of the injected frequency is $\sim 93\%$ in this case. For the DACF method we find that only $\sim 34\%$ of the simulations are recovered, although if we consider only simulations where the injected period is greater than 10 days the fraction of simulations that are recovered by DACF is then $\sim 81\%$. The fraction of these latter recoveries for which the recovered frequency is within 0.001 day^{-1} of the injected frequency is only $\sim 51\%$ in this case, however. For the simulations the recovery frequency is relatively insensitive to the period and color and depends most significantly on the amplitude. For amplitudes between 0.01 mag and 0.022 mag the recovery fraction is $\sim 65\%$, whereas for amplitudes above 0.05 mag the recovery fraction is $\sim 90\%$. Above 0.05 mag the recovery fraction is independent of amplitude. As noted above, the estimated completeness of $\sim 80\%$ is likely to be overly optimistic since we do not include the by-eye selection, do not account for the reduction in signal amplitude by TFA and use relatively “easy-to-find” sinusoid signals. Nonetheless we expect that the recovery frequency is $\gtrsim 70\%$.

As seen in figure 16 the fraction of stars that are detected as variables increases steeply with decreasing stellar mass. While only $\sim 3\%$ of stars with $M \gtrsim 0.7 M_{\odot}$ are found to be variable, approximately half of the stars with $M \lesssim 0.2 M_{\odot}$ are detected as variables with peak-to-peak amplitudes $\gtrsim 0.01$ mag (fig. 15). We find that an exponential relation of the form

$$\text{Var. Frac.} = (0.0034 \pm 0.0008)e^{(0.84 \pm 0.06)(V-K_S)} \quad (11)$$

fits the observed relation over the color range $2.0 < V - K_S < 6.0$.

In figure 17 we show the relation between period and color. For stars with $V - K_S \lesssim 4.5$ the measured distribution of $\log P$ is peaked at ~ 20 days with a broad tail toward shorter periods and a more rapid drop-off for longer periods. Note that the cut-off for longer periods may be due to the correlation between period and amplitude for these stars; stars with periods longer than ~ 20 days may be harder to detect and not intrinsically rare. The peak of the distribution appears to be correlated with color such that redder stars are found more commonly at longer periods than bluer stars. For stars with $V - K_S \gtrsim 4.5$ the distribution changes significantly such that the $\log P$ distribution appears to be more or less flat between ~ 0.3 and ~ 10 days, while red stars with $P \gtrsim 10$ days are uncommon. The morphology is broadly consistent with what has been seen from other surveys.

To demonstrate this, in Figure 18 we compare the mass-period distribution for stars in our survey to the results from other surveys of field stars and open clusters. We choose to use mass for the comparison rather than observed colors because a consistent set of colors is not available for all surveys. The masses for stars in our survey are estimated from their $V - K_S$ colors (see Fig. 16).




FIG. 16.— Top: The distribution of $V - K_S$ colors for the 1849 stars in our catalog of periodic variables that are flagged as being secure detections for at least one detection method and are not flagged as probable blends or as having problematic amplitudes, compared to the distribution of $V - K_S$ colors for all 27,560 stars in the sample. On the top axis we show the corresponding main sequence stellar masses determined by combining the empirical $V - K_S$ vs. M_K main-sequence for stars in the Solar neighborhood given by Johnson & Apps (2009) with the mass- M_K relation from the Baraffe et al. (1998) 4.5 Gyr, solar-metallicity isochrone with $L_{\text{mix}} = 1.0$. We used the relations from Carpenter (2001) to convert the CIT K magnitudes from the isochrones into the 2MASS system. The distribution for variable stars is biased toward redder $V - K_S$ colors relative to the distribution for all stars. The decrease in the total number of stars in the sample redward of $V - K_S \sim 3.5$ is due to the V -band magnitude limit of the PPMX survey. Bottom: The completeness corrected fraction of stars that are variable with peak-to-peak R or I_C amplitude > 0.01 mag as a function of $V - K_S$; this fraction increases exponentially with color (solid line, eq. 11). While only $\sim 3\%$ of stars with $M \gtrsim 0.7 M_{\odot}$ are found to be variable at the $\gtrsim 1\%$ level, approximately half of the stars with $M \lesssim 0.2 M_{\odot}$ are variable at this level.

We take data from the Mount Wilson and Vienna-KPNO (Strassmeier et al. 2000) samples of field stars with rotation periods. For the Mount Wilson sample we use the compilation by Barnes (2007), the original data comes from Baliunas et al. (1996) and from Noyes et al. (1984). For the Vienna-KPNO sample we only consider stars which are listed as luminosity class V. We estimate the masses for stars in these samples using the same $V - K_S$ to mass conversion that we use for our own sample. The V and K_S magnitudes for these field stars are taken from SIMBAD where available. We also compare our sample to four open clusters with ages between




FIG. 17.— Top: Period vs. $V - K_S$ for 1785 stars in our catalog of periodic variables that are flagged as being secure detections for at least one detection method, are not flagged as having incorrect period determinations for that method, and are not flagged as probable blends or as having problematic amplitudes. There appears to be a paucity of stars with $P > 10$ days and $V - K_S > 4.5$ or $V - K_S < 3$. Bottom: The distributions of periods are shown for four color bins. For the three bluest color bins there appears to be a correlation between period and color, such that the mode period is longer for redder stars. For stars with $V - K_S \gtrsim 4.5$, on the other hand, the period distribution appears to be biased to shorter periods. Using a K-S test, we find that the probability that the stars in any two of the different color bins are drawn from the same distribution is less than 0.01% for all combinations except the for the combination of the two intermediate color bins. For that combination the probability is $\sim 0.7\%$.

100–200 Myr, including M35 (~ 180 Myr; Meibom et al. 2009), and three clusters observed by the MONITOR project: M50 (~ 130 Myr; Irwin et al. 2009a), NGC 2516 (~ 150 Myr; Irwin et al. 2007), and M34 (~ 200 Myr; Irwin et al. 2006). Finally, we compare our sample to the two oldest open clusters with significant samples of rotation periods: M37 (~ 550 Myr; Hartman et al. 2009), and the Hyades (~ 625 Myr; Radick et al. 1987, 1995; Prosser et al. 1995). For the MONITOR clusters we use the mass estimates given in their papers, these are based on the I_C -mass relation from the appropriate Baraffe et al. (1998) isochrone for the age/metallicity of each cluster. For M35, M37 and the Hyades we use the mass estimates derived from the V, I_C -mass relations from the appropriate YREC (An et al. 2007) isochrones for each cluster. For M35 we only include 214 stars from the Meibom et al. (2009) catalog that lie near the main

sequence in V , B and I_C , and we exclude any stars which have a proper motion membership probability less than 80%, or an RV membership probability less than 80% as determined by Meibom et al. (2009). We expect that the stars in our sample, and in the other field star samples, have a range of ages, but on average will be older than the stars in the open clusters.

The sample of stars with rotation periods presented here is substantially richer than is available for other surveys of field stars. This is especially the case for later spectral types. The Mt. Wilson and Vienna-KPNO surveys primarily targeted G and early K stars, so there is not much overlap in stellar mass between those samples and our sample. The few Mt. Wilson stars with estimated masses $\lesssim 0.8 M_\odot$ do show an anti-correlation between mass and period, and have periods that are longer than the majority of stars in our sample. The Vienna-KPNO stars, on the other hand, have periods that cluster around ~ 10 days, which is closer to the mode of the period distribution for stars of comparable mass in our sample. Since the Vienna-KPNO stars were selected as showing spectroscopic evidence for chromospheric activity before their periods were measured photometrically, this sample is presumably more biased to shorter period active stars than the Mt. Wilson sample. Given the correlation between photometric amplitude and rotation period, we would also expect our sample to be biased toward shorter period stars relative to the Mt. Wilson sample. When compared to the open cluster samples we see clear evidence for evolution in the rotation periods of low-mass stars. Stars in the younger clusters have shorter periods at a given mass, on average, than stars in our sample. The discrepancy becomes more apparent for stars with $M \lesssim 0.5 M_\odot$ for which the period and mass appear to be positively correlated in the young cluster samples while they are anti-correlated in our sample. Looking at the ~ 600 Myr clusters, again the periods of stars in our sample are longer at a given mass, on average, than the periods of the cluster stars, however in this case the mode of the period distribution for the cluster stars appears to be closer to the mode of the period distribution for our sample than it is for the younger clusters. The lowest mass stars in the older clusters also do not show as significant a correlation between mass and period as do the lowest mass stars in the younger clusters. For stars with $M \lesssim 0.3 M_\odot$ the available field star and older open cluster samples are too sparse to draw any conclusions from when comparing to our sample. For the younger clusters, we note that the distribution of periods for the lowest mass stars is even more strongly peaked toward short periods than it is in our sample. This suggests that these stars do lose angular momentum over time, despite not having a tachocline. A more detailed comparison of these data to models of stellar angular momentum evolution is beyond the scope of this paper.

6.2.3. Period-X-ray Relation

Figure 19 shows the fraction of variables that match to an X-ray source as a function of period. This fraction is constant at $\sim 22\%$ for periods less than ~ 4 days, for longer periods the fraction that matches to an X-ray source decreases as $\sim P^{-0.8}$. Following Agüeros et al.




FIG. 18.— A comparison of the mass-period distribution for stars in our survey to the results from other surveys. See the text for a description of the data sources. For clarity we make the comparison separately for field stars, open clusters with $100 \text{ Myr} < t < 500 \text{ Myr}$ and for two open clusters with $t \sim 600 \text{ Myr}$. The rotation periods of stars in our sample at a given mass appear to be longer, on average, than the periods of stars in the open clusters, this is true across all mass ranges covered by our survey. The rotation periods from the Vienna-KPNO survey appear to be comparable, at a given mass, to the periods of stars in the open clusters, while the periods from the Mt. Wilson survey appear to be generally longer than the periods from our survey. It is likely that our survey and the Vienna-KPNO surveys are biased toward high-activity, shorter period stars than the Mt. Wilson survey is.

(2009) we calculate the ratio of X-ray to J-band flux via

$$\log_{10}(f_X/f_J) = \log_{10} f_X + 0.4J + 6.30 \quad (12)$$

where 1 count s^{-1} in the 0.1–2.4 keV energy range is assumed to correspond on average to $f_X = 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$. In figure 20 we plot the flux ratio as a function of rotation period for samples of variables separated by their $V - K_S$ color. The X-ray flux is anti-correlated with the rotation period for stars with $M \gtrsim 0.25 M_\odot$, for stars with $M \lesssim 0.25 M_\odot$ there is still a hint of an anti-correlation, though it is of low statistical significance (the false alarm probability is $\sim 20\%$). This result is similar to what we found for the photometric amplitude-period relation.




FIG. 19.— The fraction of non-EB periodic variable stars that match to a ROSAT source as a function of rotation period. The errorbars for the 3 longest period bins show 1σ upper-limits. For stars with rotation periods less than ~ 4 days the fraction that matches to an X-ray source is constant at $\sim 22\%$, for longer periods the fraction decreases as $\sim P^{-0.8}$.

6.3. Flares

In Section 3.4 we conducted a search for large-amplitude long-duration flares, finding only 64 events in 60 out of 23,589 stars with EPD light curves that were analyzed. There are likely to be many more flare events that have been observed but which cannot be easily distinguished from non-Gaussian noise in an automated fashion. The presence of these flares may, however, be identified statistically by looking for an excess of bright outliers relative to faint outliers in the light curves that correlates with other observables, such as the rotation period. For each light curve we determine the excess fraction of bright $n\delta$ outliers via

$$f_n = \frac{N_{n,-} - N_{n,+}}{N_{\text{tot}}} \quad (13)$$

where $N_{n,-}$ is the number of points in the light curve with $m - m_0 < -n\delta$, $N_{n,+}$ is the number of points with $m - m_0 > n\delta$, m_0 is the median of the light curve, δ is the median value of $|m - m_0|$, and there are N_{tot} points in the light curve. We do this for $n = 3, 5$ and 10 . Before calculating f_n we high-pass filter the light curve by subtracting from each point the median of all points that are within 0.1 day of that point.

It is a well known fact that the distribution of magnitudes in the light curve of a faint star is skewed about the median toward faint values. As such we expect f_n to




FIG. 20.— The ratio of 0.1-2.4 keV X-ray flux to J -band near infrared flux vs. the rotation period for non-EB periodic variable stars that match to a ROSAT source. We divide the sample into the same 4 color bins used in fig. 15. We also list the Spearman rank-order correlation coefficient and the statistical significance of the correlation for each sample. The X-ray flux is anti-correlated with the rotation period at high significance for stars with $V - K_S < 5.0$. For stars with $V - K_S > 5.0$ ($M \lesssim 0.25 M_\odot$) there is still a hint of an anti-correlation, though it is of low statistical significance.

be less than zero and to decrease for fainter stars. Figure 21 shows the excess fraction of bright outliers as a function of the light curve scatter for the high-pass filtered light curves. We compare the observed relation to the relation obtained for a simulated set of light curves generated using Poisson noise for the flux from the star, the sky and the sky annulus. We simulate one light curve for each observed light curve, using the observed sky fluxes and extinctions to determine the expected sky flux and star flux for each point in each light curve. We find that the median relation between the excess fraction of bright outliers and the light curve scatter is consistent with the expected relation; however there is greater scatter about this relation than expected from our idealized noise model. It is unclear whether other sources of noise, such as systematic variations due to blending, flat-fielding errors, pixelation effects, etc., will yield a skewed distribution of magnitude values, and if so, in which direction the magnitude distribution will be skewed. It is unlikely, however, that this would be correlated with parameters such as the rotation period. Therefore, if a correlation is observed between rotation period and the excess fraction of bright outliers, then it is likely to be physical.

In figure 22 we plot the median excess fraction of bright 5- δ and 10- δ outliers as a function of period for stars in our variables catalog that are identified as reliable detections for at least one search algorithm, and that are not flagged as probable blends or as having problematic amplitudes. In order of preference, we adopt the AoVHarm TFA, AoVHarm EPD, AoV TFA, AoV EPD, DACF TFA or DACF EPD period for the star. There appears to be a slight anti-correlation between period and excess fraction




FIG. 21.— The excess fraction of bright 5- δ outliers vs. the light curve scatter δ , shown for the $\sim 27,000$ observed light curves (top) and for a simulated sample of light curves (bottom). The dark points with the error bars show the median values of f_5 for the observed light curves, the solid line shows a power-law fit to the median values for the observed light curve, while the dashed line shows a power-law fit to the median values for the simulations.

of bright 5- δ outliers such that stars with short periods (< 10 days) have slightly more bright 5- δ outliers than faint 5- δ outliers compared to the median, while stars with long periods (> 10 days) have slightly fewer compared to the median. A Spearman rank-order correlation test rejects the null hypothesis that the excess fraction is not anti-correlated with the period at 99.92% confidence. For 10- δ outliers, the excess fraction of bright outliers also appears to be anti-correlated with the period, though at lower significance (99.5%).

In figure 22 we also compare the distribution of periods for stars with detected large-amplitude long-duration flares to the distribution for stars without such flare detections. A total of 31 of the 60 flare stars have a robust period determination and are not flagged as a probable blend. The period detection frequency of $\sim 50\%$ for flare stars is significantly higher than that for all other stars ($\lesssim 7\%$). This result is expected if stellar flaring is associated with the large starspots that give rise to continuous photometric variations. As seen in figure 22, the distribution of periods for flare stars is concentrated toward shorter periods than the distribution for non-flare stars. The longest period found for a flare star is 18.2 days whereas 31% of the non-flare stars with period determi-

nations have periods greater than 18.2 days. Conducting a K-S test we find that the probability that the two samples are drawn from the same distribution is less than 10^{-6} .

7. CONCLUSION

In this paper we have presented the results of a variability survey conducted with HATNet of field K and M dwarfs selected by color and proper motion. We used a variety of variability selection techniques to identify periodic and quasi-periodic variables, and have also conducted a search for large amplitude, long-duration flare events. We conducted Monte Carlo simulations of light curves with realistic noise properties to aid in setting the selection thresholds. Out of a total sample of 27,560 stars we selected 3496 that show potential variability, including 95 that show eclipses in their light curves, and 60 that show flares. We inspected all automatically selected light curves by eye, and flagged 2321 stars (including those with flares) as being secure variability detections. Because the HATNet images have poor spatial resolution, variability blending is a significant problem. We therefore implemented an automated routine to classify selected non-flare variables as probable blends, potential blends, unlikely blends, unblended or as having problematic amplitudes. Altogether we found 1928 variables that are classified as secure detections and are not classified as probable blends or as having problematic amplitudes (cases where the best-fit Fourier series to the light curve has a flare alarm probability greater than 10%). This includes 79 stars that show eclipses in their light curves. We identified 64 flare events in 60 stars, 38 of these stars are also selected as potential periodic or quasi-periodic variables (37 are considered reliable detections, of which 36 have reliable period determinations and 31 have reliable period determinations and are not flagged as probable blends or as having problematic amplitudes). We matched the sample of potential variables to other catalogs, and found that 77 lie within $2'$ of a previously identified variable, while 3419 do not. Including only flare stars and variables that are classified as secure detections and are not classified as probable blends or as having problematic amplitudes, 43 (including 7 EBs) lie within $2'$ of a previously identified variable, so that 1885 are new identifications.

One of the eclipsing binaries that we identified is the previously known SB2 system 1RXS J154727.5+450803. By combining the published RV curves for the component stars with the HATNet I band light curve, we obtained initial estimates for the masses and radii of the component stars (Tab. 4). The system is one of only a handful of known double-lined eclipsing binaries with component masses less than $0.3 M_{\odot}$. While we caution that the errors on the component radii are likely to be underestimated due to systematic errors that have not been considered in this preliminary analysis, it is interesting that the radii do appear to be larger than predicted if the system is older than ~ 200 Myr. With a magnitude $V \sim 13.4$, this system is only slightly fainter than the well-studied binary CM Dra ($V \sim 12.90$) which has been the anchor of the empirical mass-radius relation for very late M dwarfs. 1RXS J154727.5+450803 is thus a promising target for more detailed follow-up to obtain high precision measurements of the fundamental param-




FIG. 22.— Top: The median excess fraction of bright 5δ outliers and 10δ outliers vs. period for stars in our variables catalog that are identified as reliable detections for at least one search algorithm, and that are not flagged as probable blends or as having problematic amplitudes. In computing the excess fraction for each light curve we subtract the median excess fraction for the light curve scatter (fig. 21). Center: The median rank of the excess fraction of bright 5δ and 10δ outliers vs. the rank period. The anti-correlation between period and the excess fraction of bright outliers is more apparent when the ranks, rather than the values, are plotted against each other. Bottom: Comparison between the period distributions for stars with detected large-amplitude long-duration flares, and stars without such a flare detected. Note the absence of long-period stars with flares. For clarity we have multiplied the flare star period distribution by a factor of 10.

eters of the component stars. With additional follow-up,




FIG. 23.— Top: The distribution of $V - K_S$ colors for the 60 stars with high amplitude flares detected compared to the distribution of $V - K_S$ colors for all 27,560 stars in the sample. The distribution for flares stars is biased toward redder $V - K_S$ colors relative to the distribution for all stars. Note that we use a 5 times higher binning resolution for the full sample. Bottom: The fraction of stars with a high amplitude flare detection is plotted against $V - K_S$. For stars with $V - K_S < 4.0$ less than $\sim 0.1\%$ of stars had a flare detected, for stars with $V - K_S > 4.5$ the fraction is $\gtrsim 1\%$.

the large sample of 79 probable late-type eclipsing binaries presented in this paper should prove fruitful for further investigations of the fundamental parameters of low-mass stars.

The majority of the variable stars that we have identified are likely to be BY Dra type variables, with the measured period corresponding to the rotation period of the star. This is the largest sample of rotation periods presented to date for late-type field stars. We discussed a number of broad trends seen in the data, including an anti-correlation between the rotation period and the photometric amplitude of variability, an exponential relation between $V - K_S$ color and the fraction of stars that are variable, a positive correlation between period and the $V - K_S$ color for stars with $V - K_S \lesssim 4.5$, a relative absence of stars with $P \gtrsim 10.0$ days and $V - K_S \gtrsim 4.5$, and an anti-correlation between the rotation period and the ratio of X-ray to J band flux. The correlations between period and activity indicators including the amplitude of photometric variability and the X-ray emission are consistent with the well-known rotation-age-activity-mass relations for F, G, K and early M dwarfs. The

data presented here may help in further refining these relations. Our data hints at a change in the rotation-activity connection for the least massive stars in the sample ($M \lesssim 0.25 M_\odot$). The anti-correlation between period and amplitude appears to break down for these stars, and similarly the period-X-ray anti-correlation is less significant for these stars than for more massive stars. This is potentially at odds with previous studies which used $H\alpha$ to trace activity and $v \sin i$ to infer rotation period, and found that the period-activity anti-correlation extends to very late-type M dwarfs. Comparing our sample to other field and open cluster samples, we find that the rotation periods of stars in our sample are generally longer than the periods found in open clusters with $t \lesssim 620$ Myr, which implies that K and M dwarf stars continue to lose angular momentum past the age of the Hyades. This appears to be true as well for stars with $M \lesssim 0.25 M_\odot$, though these stars generally have shorter periods than more massive stars in our sample.

Finally we have conducted a search for flare events in our light curves, identifying 64 events in 60 stars. Due to the difficulty between distinguishing a flare from bad photometry in an automated way, there are likely to be many flare events in the light curves that we do not identify. We therefore do not attempt to draw conclusions about the total occurrence rate of flaring (for a recent determination of this frequency using data from the Sloan Digital Sky Survey, see Kowalski et al. 2009). We find that the distribution of $V - K_S$ colors for flare stars is biased toward red colors, implying that the flare frequency increases with decreasing stellar mass, which has been known for a long time (Ambartsumyan et al. 1970, see also Kowalski et al. 2009). We find that roughly half the flare stars are detected as periodic variables, which is a significantly higher fraction than for the full sample of stars. This is in line with the expectation that stellar flaring is associated with the presence of significant starspots, and is consistent with the finding by Kowalski et al. (2009) that the flaring frequency of active M dwarfs showing $H\alpha$ emission is ~ 30 times higher than the flaring frequency of inactive M dwarfs. We also find that the distribution of periods for flare stars is biased toward shorter periods, again as expected from the rotation-activity connection. Finally we attempt to statistically identify flares by searching for excess bright outliers relative to faint outliers in the light curves. This excess appears to be anti-correlated with rotation period and provides further evidence that flares are more common among rapidly rotating K and M dwarfs than among slower rotators.

HATNet operations have been funded by NASA grants NNG04GN74G, NNX08AF23G and SAO IR&D grants. G.Á.B. acknowledges support from the Postdoctoral Fellowship of the NSF Astronomy and Astrophysics Program (AST-0702843). T.M. acknowledges support from the ISRAEL SCIENCE FOUNDATION (grant No. 655/07). The Digitized Sky Surveys were produced at the Space Telescope Science Institute under U.S. Government grant NAG W-2166. The images of these surveys are based on photographic data obtained using the Oschin Schmidt Telescope on Palomar Mountain and the UK Schmidt Telescope. The plates were processed into

the present compressed digital form with the permission of these institutions. This research has made use of data obtained from or software provided by the US National Virtual Observatory, which is sponsored by the

National Science Foundation. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France.

REFERENCES

- Agüeros, M. A., et al. 2009, *ApJS*, 181, 444
 Aigrain, S., et al. 2008, *A&A*, 488, L43
 Akerlof, C., et al. 2000, *AJ*, 119, 1901
 An, D., Terndrup, D. M., Pinsonneault, M. H., Paulson, D. B., Hanson, R. B., & Stauffer, J. R. 2007, *ApJ*, 655, 233
 Ambartsumyan, V. A., Mirzoyan, L. V., Parsamyan, E. S., Chavushyan, O. S., & Erastova, L. K. 1970, *Astrofizika*, 6, 7
 Bakos, G., Noyes, R. W., Kovács, G., Stanek, K. Z., Sasselov, D. D., & Domsa, I. 2004, *PASP*, 116, 266
 Bakos, G. Á., et al. 2009, *ApJ*, in press, arXiv:0901.0282
 Baliunas, S. L., et al. 1995, *ApJ*, 438, 269
 Baliunas, S., Sokoloff, D., & Soon, W. 1996, *ApJ*, 457, L99
 Baraffe, I., Chabrier, G., Allard, F., & Hauschildt, P. H. 1998, *A&A*, 337, 403
 Barnes, S. A. 2007, *ApJ*, 669, 1167
 Beatty, T. G., et al. 2007, *ApJ*, 663, 573
 Bessell, M. S., & Brett, J. M. 1988, *PASP*, 100, 1134
 Blake, C. H., Torres, G., Bloom, J. S., & Gaudi, B. S. 2008, *ApJ*, 684, 635
 Bonatto, Ch., Bica, E., & Girardi, L. 2004, *A&A*, 415, 571
 Carpenter, J. M. 2001, *AJ*, 121, 2851
 Chabrier, G., & Baraffe, I. 1997, *A&A*, 327, 1039
 Chabrier, G., Baraffe, I., Allard, F., & Hauschildt, P. 2000, *ApJ*, 542, 464
 Chabrier, G., & Küker, M. 2006, *A&A*, 446, 1027
 Chabrier, G., Gallardo, J., & Baraffe, I. 2007, *A&A*, 472, L17
 Claret, A. 2000, *A&A*, 363, 1081
 Cooley, J. W., & Tukey, J. W. 1965, *Mathematics of Computation*, 19, 297
 Cousins, A. W. J. 1976, *MmRAS*, 81, 25
 Creevey, O. L., et al. 2005, *ApJ*, 625, 127
 Delfosse, X., Forveille, T., Perrier, C., & Mayor, M. 1998, *A&A*, 331, 581
 Demory, B.-O., et al. 2009, *A&A*, in press, arXiv:0906.0602v1
 Devor, J. 2005, *ApJ*, 628, 411
 Devor, J., et al. 2008, *ApJ*, 687, 1253
 Duncan, D. K., et al. 1991, *ApJS*, 76, 383
 Edelson, R. A., & Krolik, J. H. 1988, *ApJ*, 333, 646
 Elias, J. H., Frogel, J. A., Hyland, A. R., & Jones, T. J. 1983, *AJ*, 88, 1027
 Elias, J. H., Frogel, J. A., Matthews, K., & Neugubauer, G. 1982, *AJ*, 87, 1029
 Etzel, P. B. 1981, *NATO ASI*, p. 111
 Fernandez, J., et al. 2009, *ApJ*, in press, arXiv:0906.2207
 Gould, A., & Morgan, C. W. 2003, *ApJ*, 585, 1056
 Guinan, E. F., & Engle, S. G. 2009, arXiv:0901.1860
 Hartman, J. D., Bakos, G., Stanek, K. Z., & Noyes, R. W. 2004, *AJ*, 128, 1761
 Hartman, J. D., Gaudi, B. S., Holman, M. J., McLeod, B. A., Stanek, K. Z., Barranco, J. A., Pinsonneault, M. H., & Kalirai, J. S. 2008, *ApJ*, 675, 1254
 Hartman, J. D., et al. 2009, *ApJ*, 691, 342
 Hebb, L., Wyse, R. F. G., Gilmore, G., & Holtzman, J. 2006, *AJ*, 131, 555
 Høg, E., et al. 2000, *A&A*, 355, L27
 Horne, J. H., & Baliunas, S. L. 1986, *ApJ*, 302, 757
 Irwin, J., Aigrain, S., Hodgkin, S., Irwin, M., Bouvier, J., Clarke, C., Hebb, L., & Moraux, E. 2006, *MNRAS*, 370, 954
 Irwin, J., Hodgkin, S., Aigrain, S., Hebb, L., Bouvier, J., Clarke, C., Moraux, E., & Bramich, D. M. 2007, *MNRAS*, 377, 741
 Irwin, J., Aigrain, S., Bouvier, J., Hebb, L., Hodgkin, S., Irwin, M., & Moraux, E. 2009a, *MNRAS*, 392, 1456
 Irwin, J., et al. 2009b, *ApJ*, in press, arXiv:0906.4365
 Johnson, J. A., & Apps, K. 2009, *ApJ*, in press, arXiv:0904.3092
 Karoff, C., et al. 2007, *AJ*, 134, 766
 Kasting, J. F., Whitmire, D. P., & Reynolds, R. T. 1993, *Icarus*, 101, 108
 Kazarovets, E. V., Samus', N. N., & Durevich, O. V. 1998, *IBVS*, 4655
 Kharchenko, N. V. 2001, *Kinematika i Fizika Nebesnykh Tel*, 17, 409
 Kholopov, P. N. (ed) 1982, *The New Catalogue of Suspected Variable Stars*, Moscow: Nauka
 Kovács, G. 1980, *Ap&SS*, 69, 485
 Kovács, G., Zucker, S., & Mazeh, T. 2002, *A&A*, 391, 369
 Kovács, G., Bakos, G., & Noyes, R. W. 2005, *MNRAS*, 356, 557
 Kowalski, A. F., Hawley, S. L., Hilton, E. J., Becker, A. C., West, A. A., Bochanski, J. J., & Sesar, B. 2009, arXiv:0906.2030
 Krautter, J. 1996, in *Light Curves of Variable Stars: a Pictorial Atlas*, eds. C. Sterken and C. Jaschek (Cambridge: Cambridge University Press), 53
 Lacy, C. H. 1977, *ApJ*, 218, 444
 Landolt, A. U. 1992, *AJ*, 104, 340
 Lammer, H., et al. 2007, *Astrobiology*, 7, 185
 Lomb, N. R. 1976, *A&SS*, 39, 447
 López-Morales, M. 2005, *ApJ*, 631, L1120
 López-Morales, M. 2007, *ApJ*, 660, L732
 Lupton, R. 1993, *Statistics in Theory and Practice*, (Princeton: Princeton University Press)
 Luyten, W. J. 1922, *Lick Observatory Bulletin*, 10, 135
 Mamajek, E. E., & Hillenbrand, L. A. 2008, *ApJ*, 687, 1264
 Marigo, P., Girardi, L., Bressan, A., Groenewegen, M. A. T., Silva, L., & Granato, G. L. 2008, *A&A*, 482, 883
 Marquardt, D. W. 1963, *Journal of the Society for Industrial and Applied Mathematics*, 11, 431
 Mason, K. O., et al. 2000, *MNRAS*, 311, 456
 Maxted, P. F. L., O'Donoghue, D., Morales-Rueda, L., Napiwotzki, R., & Smalley, B. 2007, *MNRAS*, 376, 919
 Meibom, S., Mathieu, R. D., & Stassun, K. G. 2009, *ApJ*, 695, 679
 Messina, S., Rodonò, M., & Guinan, E. F. 2001, *A&A*, 366, 215
 Metcalfe, T. S., Mathieu, R. D., Latham, D. W., & Torres, G. 1996, *ApJ*, 456, 356
 Mirzoyan, L. V., Ambaryan, V. V., Garibdzhanyan, A. T., & Mirzoyan, A. L. 1989, *Astrophysics*, 31, 567
 Mochnacki, S. W., et al. 2002, *AJ*, 124, 2868
 Mohanty, S., & Basri, G. 2003, *ApJ*, 583, 451
 Monet, D. G., et al. 2003, *AJ*, 125, 984
 Morales, J. C., et al. 2009, *ApJ*, 691, 1400
 Nataf, D. M., Stanek, K. Z., & Bakos, G. A. 2009, *AcA* submitted, arXiv:0904.4696
 Nelson, B., & Davis, W. D. 1972, *ApJ*, 174, 617
 Norton, A. J., et al. 2007, *A&A*, 467, 785
 Noyes, R. W., Hartmann, L., Baliunas, S. L., Duncan, D. K., & Vaughn, A. H. 1984, *ApJ*, 279, 763
 Paczyński, B. 1997, "The Future of Massive Variability Searches," in *Proceedings 12th IAP Colloquium: "Variable Stars and the Astrophysical Returns of Microlensing Searches"*, Paris (Ed. R. Ferlet), 357
 Pál, A., & Bakos, G. Á. 2006, *PASP*, 118, 1474
 Pál, A. 2009, Ph.D. Thesis, Eötvös Loránd University, arXiv:0906.3486
 Parley, N. R., et al. 2006, *Earth, Moon, and Planets*, 97, 261
 Perryman, M. A. C., et al. 1997, *A&A*, 323, L49
 Pizzolato, N., Maggio, A., Micela, G., Sciortino, S., & Ventura, P. 2003, *A&A*, 397, 147
 Pojmanski, G. 2002, *AcA*, 52, 397
 Pont, F., Zucker, S., & Queloz, D. 2006, *MNRAS*, 373, 231
 Popper, D. M., & Etzel, P. B. 1981, *AJ*, 86, 102
 Press, W. H., & Rybicki, G. B. 1989, *ApJ*, 338, 277
 Press, W. H., Teukolsky, S. A., Vetterling, W. T., & Flannery, B. P. 1992, *Numerical Recipes in C*, 2nd ed. (New York: Cambridge University Press)
 Prosser, C. F., et al. 1995, *PASP*, 107, 211
 Radick, R. R., Thompson, D. T., Lockwood, G. W., Duncan, D. K., & Baggett, W. E. 1987, *ApJ*, 321, 459
 Radick, R. R., Lockwood, G. W., Skiff, B. A., & Thompson, D. T. 1995, *ApJ*, 452, 332
 Reiners, A., & Basri, G. 2007, *ApJ*, 656, 1121
 Reyle, C., & Robin, A. C. 2004, *A&A*, 421, 643
 Ribas, I. 2003, *A&A*, 398, 239
 Ribas, I. 2006, *Ap&SS*, 304, 89

- Röser, S., Schilbach, E., Schwan, H., Kharchenko, N. V., Piskunov, A. E., & Scholz, R.-D. 2008, *A&A*, 488, 401
- Samus', N. N., Durlevich, O. V., Zharova, A. V., Kazarovets, E. V., Kireeva, N. N., Pastukhova, E. N., Williams, D. B., & Hazen, M. L. 2006, *Astronomy Letters*, 32, 263
- Scargle, J. D. 1982, *ApJ*, 263, 835
- Schwarzenberg-Czerny, A. 1989, *MNRAS*, 241, 153
- Schwarzenberg-Czerny, A. 1996, *ApJ*, 460, L107
- Shporer, A., Mazeh, T., Moran, A., Bakos, G., & Kovács, G. 2007, in "Transiting Extrasolar Planets Workshop, ASP Conference Series 366" eds. C. Afonso, D. Weldrake, & Th. Henning, 99
- Shporer, A., Bakos, G. Á., Mazeh, T., Kovács, G., & Sipőcz, B. 2009, *IAU Symposium*, 253, 331
- Skrutskie, M. F., et al. 2006, *AJ*, 131, 1163
- Skumanich, A. 1972, *ApJ*, 171, 565
- Southworth, J., Maxted, P. F. L., & Smalley, B. 2004, *MNRAS*, 351, 1277
- Southworth, J., Zucker, S., Maxted, P. F. L., & Smalley, B. 2004, *MNRAS*, 355, 986
- Southworth, J., Smalley, B., Maxted, P. F. L., Claret, A., & Etzel, P. B. 2005, *MNRAS*, 363, 529
- Stellingwerf, R. F. 1978, *ApJ*, 224, 953
- Strassmeier, K., Washuettl, A., Granzer, T., Scheck, M., & Weber, M. 2000, *A&AS*, 142, 275
- Szulágyi, J., Kovács, G., & Welch, D. L. 2009, *A&A*, in press, arXiv:0903.1165
- Torres, G., & Ribas, I. 2002, *ApJ*, 567, 1140
- Torres, G., Lacy, C. H., Marschall, L. A., Sheets, H. A., & Mader, J. A. 2006, *ApJ*, 640, 1018
- Torres, G. 2007, *ApJ*, 671, L65
- van den Besselaar, E. J. M., et al. 2007, *A&A*, 466, 1031
- Voges, W., et al. 1999, *A&A*, 349, 389
- West, A. A., & Basri, G. 2009, *ApJ*, 693, 1283
- Wilson, O. C., & Bappu, M. K. V. 1957, *ApJ*, 125, 661
- Wilson, O. C. 1978, *ApJ*, 226, 379
- Yong, D., & Lambert, D. L. 2003, *PASP*, 115, 796

APPENDIX

A. MONTE CARLO SIMULATIONS TO DETERMINE SELECTION CRITERIA

Here we describe the Monte Carlo simulations that we conducted to inform our choice of selection criteria for several of the variability detection algorithms that we discussed in § 3. The goal of these simulations is to establish thresholds on each variability statistic such that one may be fairly confident that light curves with values above the threshold exhibit real variability. Under the ideal case where the light curves consist of uniformly sampled Gaussian white noise, the probability distribution for many of the variability statistics may be calculated analytically, and a formal false alarm probability may be determined. For nonuniform time sampling, however, the analytic expressions generally do not hold. In this case one may still use an analytic approximation for the false alarm probability by including a parameter for the effective number of independent trial frequencies. This parameter must either be determined by conducting Monte Carlo simulations (e.g. Horne & Baliunas 1986), or it can be estimated from the total number of data points and the total number of frequencies sampled (see Press et al. 1992). In general, however, the noise in a light curve is neither Gaussian nor uncorrelated in time. Temporally-correlated, low-frequency ("red") noise appears to be ubiquitous in ground based variability surveys (e.g. Pont, Zucker & Queloz 2006); additionally the distribution of magnitude values in ground based light curves often show broad non-Gaussian wings. As we will show below, the HATNet light curves exhibit both of these effects. In both cases the analytic expressions derived for Gaussian white noise will underestimate the actual false alarm probability. In this case one must conduct Monte Carlo simulations of light curves with realistic noise properties to establish the false alarm probabilities.

The formally proper way to proceed would be to conduct many thousand simulations for *each* real light curve to empirically determine the distribution of AoV, AoVHarm, etc. values at each frequency in the periodogram of the light curve. These distributions would then be used in selecting the peak frequency (greatest outlier) in the periodogram, and to determine the false alarm probability of finding that peak. Such a procedure is unpractical, however, for more than a few tens of stars given the non-negligible amount of time required to calculate periodograms for a single simulated light curve (calculating the AoVHarm periodogram for a typical HATNet light curve takes ~ 2 minutes running as a single process on a 2.5 GHz AMD 9850 Quad-Core Processor). Instead we opt for a simpler procedure of generating an ensemble of light curves meant to represent the observed sample of light curves if they consisted only of noise. We run the variability selection routines on the ensemble and compare the resulting period vs. S/N etc. distributions to the real distributions. Selection criteria are adopted such that few of the light curves in the simulated sample would pass the cuts. Because the noise properties of the light curves do depend on several parameters, we divide the full sample of light curves into several sub-samples and establish separate criteria for each subsample as described below.

In general the noise properties of a light curve will depend on many factors including the brightness of the star, its proximity to other stars, and its location on the image, to name just a few. Exploring all of these factors, however, is not practical given the computational constraints. Here we make the assumption that the noise depends only on the light curve type (EPD or TFA), the chip format (2K or 4K CCDs), and the magnitude of the star. For each chip format and filtering method we divide the sample of light curves into magnitude bins. We use a binsize of 1.0 mag and ranges from $6.0 < I < 14.0$ for the 2K CCDs and $7.0 < R < 15.0$ for the 4K CCDs. For the moment we exclude composite light curves that include both 2K and 4K observations. We conduct the simulations separately for each bin.

We discuss first how we generate the simulated light curves, we then discuss how the noise properties are determined for each sub-sample of light curves. To simulate a light curve with red noise we use the following procedure:

1. Choose an observed light curve as a basis for the time sampling of the simulated light curve (the simulated light curve will be interpolated to this time basis at the last step).
2. Generate a series of uncorrelated random magnitude measurements drawn from an empirical distribution of magnitudes $\hat{\rho}(m)$. We discuss below how this empirical distribution is determined. We will refer to this series as $s(t)$. The total number of generated points N_s should be a power of 2 (i.e. $\log_2 N_s$ is an integer) and such

that $N_S \Delta > T$ where Δ is the minimum time separation between points in the observed light curve and T is the total time-span covered by the observed light curve.

3. Use the fast Fourier transform (FFT) algorithm (e.g. Cooley & Tukey 1965) to calculate $\tilde{s}(\nu)$, the discrete Fourier transform (DFT) of $s(t)$.
4. To add red noise to the simulated signal we apply the filter $H(\nu)$ to $\tilde{s}(\nu)$ to get $\tilde{r}(\nu)$

$$\tilde{r}(\nu) = H(\nu)\tilde{s}(\nu). \quad (\text{A1})$$

The filter $H(\nu)$ satisfies

$$H(\nu) = \sqrt{P(\nu)} \quad (\text{A2})$$

where $P(\nu)$ is the power spectrum of the red noise. We discuss below how it is determined. Note that we assume $H(\nu)$ is real valued.

5. Apply the inverse FFT to $\tilde{r}(\nu)$ to obtain $r(t)$, a uniformly sampled time-series with the desired noise properties.
6. Interpolate $r(t)$ at the time values of the observed light curve to yield the simulated light curve in its final form. Note that because the variability selection algorithms are insensitive to scale changes in the magnitudes, we do not need to adjust the RMS of the simulated light curve to equal that of an observed light curve.

To determine the distribution of magnitude values we divide each light curve by its standard deviation calculated with 5σ iterative clipping about the median. The normalized magnitude values from all the light curves in a sub-sample are then grouped into a single list, which we take as $\hat{\rho}(m)$ in step 2 above. Note that we do not attempt to remove variable stars, or to whiten the light curves before generating this empirical distribution. Figure 24 shows the histogram of normalized magnitude values for 2K, EPD light curves with $9.0 < I < 10.0$. Note that the wings of the distribution are non-Gaussian. We find that a function of the form

$$\rho(m) = A \exp(-m^2/2\sigma_m^2) + B \exp(-|m|^{0.5}/C) \quad (\text{A3})$$

fits the observed distribution, with $\sigma_m = 0.886$, $C = 0.427$ and $A/B = 1.21$ for the aforementioned sub-sample. In simulating the light curves we draw points from the empirical distribution $\hat{\rho}(m)$, we give the functional form of the distribution for reference only. As a test on our assumption that the wings of the distribution are not dominated by contributions from variable stars or stars with significant trends, we also computed the distribution for the sub-sample shown in figure 24 after excluding stars identified as potential variables in § 3 and median filtering the light curves (subtracting from each point in the light curve the median value of all points within 0.1 day of the given point); the differences between the distributions are negligible.

To determine the power spectrum of the red noise, $P(\nu)$, used in eq. A2 we calculate the DFT power spectra for 100 randomly selected light curves in each sub-sample. The power spectra are interpolated to a common frequency grid and scaled to have unit median. These power spectra are then median combined to yield the median power spectrum $P_{\text{med}}(\nu)$. Figure 25 shows examples of $P_{\text{med}}(\nu)$ for several of the light curve sub-samples. Note the general rise in the power spectra at lower frequencies as well as the excess power near the sidereal frequency $\nu_{\text{sid}} = 1.0027379 \text{ day}^{-1}$ and its harmonics. At exactly ν_{sid} the power spectra drop to low values due to the decorrelations in the EPD routine. Also note that the power spectra of fainter stars, and of the TFA light curves show less frequency dependence (i.e. they are closer to a horizontal line, which is what one would see for white noise). As shown in figure 25, we fit a function of the form

$$P(\nu) = \frac{a/(1 + (\nu/b)^c) + d}{1 + e|\sin(\frac{\nu}{\nu_{\text{sid}}}\pi)|} + f \quad (\text{A4})$$

to $P_{\text{med}}(\nu)$, where a , b , c , d , e and f are fitting parameters which are listed for each sub-sample in table 6. The parameters a and e tend to decrease with increasing magnitude, and are generally smaller for the TFA sub-samples than for the EPD sub-samples. Otherwise, there are no clear trends in the parameter values. We use this analytic expression for $P(\nu)$ in simulating the light curves (eq. A2).

Figure 26 shows examples of several simulated light curves together with examples of some real light curves that are not selected as variable. The general noise properties of the observed light curves, including an abundance of outliers, and the presence of red noise, are reproduced in the simulated light curves.

It is important to note that there are some properties of the noise in the real light curves that are not modeled in the simulated light curves. In the real light curves the scatter of the magnitude measurements about the expected value varies in time. On nights with poor weather conditions the scatter is often higher than on photometric nights. In our simulated light curves the scatter is constant though the expected value may vary in time. Also note that if the red noise properties of the light curves vary significantly within a given sample, it is likely the median power spectrum will show less red noise than any given light curve. For this reason we adopt conservative selection criteria that pass well above the simulated variability distributions in the selections described in § 3.




FIG. 24.— The histogram of normalized Δ magnitude values for 2K, EPD light curves with $9.0 < I < 10.0$ (light filled circles). The light curves that contribute to this distribution have been divided by their standard deviations calculated with 5σ iterative clipping about the median. The solid line shows a fit to the observed distribution (eq A3). Note that we use a logarithmic scale for the vertical axis to highlight the wings of the distribution which are non-Gaussian.




FIG. 25.— Median DFT power spectra for 4 representative sub-samples of light curves. In all cases there is a rise in power toward lower frequencies which is symptomatic of red noise. Light curves also generally show increased power near the sidereal frequency and its harmonics. At exactly the sidereal frequency the power drops due to the decorrelation against variations at this frequency when applying EPD. The red noise is less significant for fainter stars, and it is less significant after trend filtering with TFA (note that this is general TFA procedure, not the signal-reconstruction mode). That some red noise remains in the post-TFA light curves suggests that for many of the fields the TFA procedure was run with fewer than the ideal number of templates required to achieve optimum noise filtering (Kovács, Bakos & Noyes 2005; Szulágyi, Kovács & Welch 2009). The solid lines show fits to the observed power spectra using eq. A4. These fits are used to simulate light curves with red noise. Note that power is plotted on a logarithmic scale.

TABLE 6
PARAMETERS FROM FITTING EQ. A4 TO THE MEDIAN POWER SPECTRA OF
DIFFERENT LIGHT CURVE SUB-SAMPLES.

Chip	LC Type	Mag.	a	b	c	d	e	f
2K	EPD	6.5	12.466	1.897	2.190	0.179	2.338	0.281
2K	EPD	7.5	21.315	1.954	2.464	0.487	5.235	0.273
2K	EPD	8.5	27.054	1.910	2.194	0.875	8.467	0.195
2K	EPD	9.5	13.549	2.054	2.319	1.114	4.635	0.136
2K	EPD	10.5	8.118	2.184	1.957	0.265	4.009	0.437
2K	EPD	11.5	6.415	2.488	2.043	0.190	4.177	0.508
2K	EPD	12.5	5.962	2.540	2.270	0.258	4.630	0.584
2K	EPD	13.5	2.862	2.842	2.073	0.191	1.882	0.557
2K	TFA	6.5	2.327	2.899	2.375	0.600	1.343	0.336
2K	TFA	7.5	2.704	2.875	2.091	0.713	1.252	0.184
2K	TFA	8.5	4.242	2.683	2.163	1.001	1.758	0.030
2K	TFA	9.5	1.530	2.793	2.181	0.962	0.539	0.000
2K	TFA	10.5	1.095	2.794	2.173	0.933	0.305	0.003
2K	TFA	11.5	0.645	3.315	2.357	0.899	0.107	0.000
2K	TFA	12.5	0.372	3.763	2.459	0.949	0.050	0.000
2K	TFA	13.5	0.296	3.828	2.472	0.958	0.027	0.000
4K	EPD	7.5	7.088	2.454	2.168	0.001	1.075	0.281
4K	EPD	8.5	8.024	2.483	2.438	0.006	1.733	0.385
4K	EPD	9.5	17.608	2.211	2.735	0.915	6.401	0.297
4K	EPD	10.5	21.052	2.081	2.515	0.767	6.782	0.254
4K	EPD	11.5	7.768	2.361	2.519	0.415	5.382	0.570
4K	EPD	12.5	6.790	2.257	2.191	0.167	5.061	0.641
4K	EPD	13.5	3.331	2.748	2.358	0.246	3.157	0.663
4K	EPD	14.5	1.794	3.330	2.239	0.121	2.188	0.738
4K	TFA	7.5	1.723	3.507	2.514	0.022	1.836	0.540
4K	TFA	8.5	1.848	3.459	2.358	0.767	0.574	0.064
4K	TFA	9.5	3.031	2.868	2.211	0.676	0.949	0.132
4K	TFA	10.5	2.226	2.670	2.158	0.791	0.419	0.000
4K	TFA	11.5	0.723	3.361	2.229	0.909	0.125	0.000
4K	TFA	12.5	0.307	3.857	2.694	0.998	0.047	0.000
4K	TFA	13.5	0.182	3.908	2.735	1.041	0.040	0.001
4K	TFA	14.5	0.118	4.340	2.896	1.084	0.069	0.000

B. CATALOG OF VARIABLE STARS

We combine the results from the various selection algorithms to form 3 catalogs of variable stars: those selected by the AoV, AoVHarm or DACF methods applied to either the EPD or TFA light curves and not flagged as an EB during the visual inspection for that method, a catalog of stars with eclipses in their light curves, and a catalog of flares. All three catalogs will be made available in electronic form from the CDS archive server¹⁰ as well as from the HATNet website¹¹. Light curves for the variable stars will be made available through the Harvard University Time Series Center¹² and through NStED¹³. There are a total of 3418 stars that are in the first catalog, 95 candidate EB stars in the second, and 64 flare events from 60 stars in the third. Note that some of the EBs and flare stars are included in the primary catalog as well. Altogether 3496 stars are included in at least one of the catalogs.

In the primary catalog of periodic variables we include the internal HAT-ID of each source, its J2000 position from 2MASS, the periods, S/N, peak-to-peak amplitudes and quality flags for the AoV and AoVHarm algorithms applied to the TFA and EPD light curves, the periods, period uncertainties, DACF peak heights, false alarm probabilities, peak-to-peak amplitude and quality flags for the DACF algorithm applied to the TFA and EPD light curves, and the blending flag described in § 4. For convenience we also include the PPMX proper motion, the 2MASS photometry, and the estimated V magnitude for each source. For each selection method we list the peak-to-peak amplitude determined for both the EPD and TFA light curves phased at that period. See § 4 for a description of how the amplitude is measured. Note that the TFA light curves used for the amplitude determination have not had been processed with signal-reconstruction mode TFA, as a result these amplitudes are generally lower than the real amplitude of the star, and are lower than the amplitudes measured on the EPD light curves. Tables 7-11 at the end of the paper show the first ten entries in this catalog. For display purposes we split the catalog into 5 separate tables, in the full electronic version of the catalog the tables are joined.

In the catalog of candidate eclipsing binary stars we include the HAT-ID, position, proper motion, 2MASS photometry, and V magnitude, together with the period, period uncertainty, epochs of primary and secondary eclipse (if the secondary is detected) and their respective uncertainties, the durations of the primary and secondary eclipses and the blending flag. Tables 12-13 at the end of the paper show the first ten entries in this catalog.

In the catalog of flare events we include the HAT-ID, position, proper motion, 2MASS photometry, and V magnitude,

¹⁰ <http://cdsarc.u-strasbg.fr>

¹¹ <http://hatnet.hu/>

¹² <http://timemachine.iic.harvard.edu/>

¹³ <http://nsted.ipac.caltech.edu>




FIG. 26.— Examples of simulated light curves (right column) compared with the observed light curves used as bases for the time sampling (left column). We show an example with significant red noise (top row), and one with very little red noise (bottom row). The observed light curves have been normalized to have unit standard deviation after an iterative 5σ clipping.

together with the time of the flare peak, the peak intensity relative to the non-flaring intensity (A in Eq. 7), and the decay time of the flare (τ in Eq. 7). Tables 14-15 at the end of the paper show the first ten entries in this catalog.

TABLE 7
CATALOG OF CANDIDATE PERIODIC VARIABLE STARS. I: ASTROMETRIC AND PHOTOMETRIC DATA

ID	RA ^a h:m:s	DE ^a deg:m:s	pmRA ^b mas/yr	pmDE ^b mas/yr	Jmag ^c mag	Hmag ^c mag	Kmag ^c mag	Vmag mag	Vref ^d	Bl ^e
HAT-085-0004278	23:22:00.71	+56:59:19.9	392.08	74.00	9.473	8.868	8.655	13.237	0	2
HAT-085-0053487	23:21:24.35	+56:25:14.3	110.04	-73.81	12.350	11.679	11.566	14.636	2	2
HAT-086-0001701	00:13:38.42	+52:45:05.0	35.13	-36.72	9.519	8.763	8.658	12.136	1	1
HAT-086-0002357	23:51:38.51	+49:45:21.3	324.91	290.31	9.062	8.400	8.210	11.437	0	1
HAT-086-0003825	23:54:10.49	+51:41:09.9	249.11	51.86	9.574	9.000	8.745	13.139	1	2
HAT-086-0004017	23:54:23.85	+54:41:26.9	35.30	35.51	9.565	8.804	8.643	12.235	1	2
HAT-086-0004248	00:07:19.34	+52:41:31.6	325.06	-3.32	9.736	9.157	8.940	12.568	0	2
HAT-086-0005153	23:51:18.20	+54:35:16.2	76.13	-7.77	9.906	9.239	9.060	12.822	1	2
HAT-086-0005542	00:16:56.01	+50:37:49.0	146.68	-27.87	9.980	9.349	9.137	12.767	1	2
HAT-086-0005602	00:16:01.61	+48:55:56.9	-83.81	51.47	10.008	9.409	9.186	12.870	1	3

^a J2000, from 2MASS^b from PPMX^c from 2MASS^d Source for Vmag (0: PPMX (Tycho-2), 1: Transformed from USNO-B1.0, 2: Transformed from 2MASS)^e Blending Flag (0: unblended, 1: unlikely blend, 2: possible blend, 3: probable blend, 4: problematic)

TABLE 8
CATALOG OF CANDIDATE PERIODIC VARIABLE STARS. II: AoV

ID	P _{ae} ^a d	SN _{ae} ^a	Amp _{ae} ^b mag	Amp _{paet} ^b mag	Q _{ae} ^{a,c}	P _{at} ^a d	SN _{at} ^a	Amp _{ate} ^b mag	Amp _{att} ^b mag	Q _{at} ^{a,c}
HAT-085-0004278	-99.999999	-999.99	-9.999	-9.999	2	-99.999999	-999.99	-9.999	-9.999	2
HAT-085-0053487	-99.999999	-999.99	-9.999	-9.999	2	14.172864	26.52	0.412	0.202	1
HAT-086-0001701	-99.999999	-999.99	-9.999	-9.999	2	18.173471	24.28	0.010	0.006	0
HAT-086-0002357	-99.999999	-999.99	-9.999	-9.999	2	-99.999999	-999.99	-9.999	-9.999	2
HAT-086-0003825	-99.999999	-999.99	-9.999	-9.999	2	-99.999999	-999.99	-9.999	-9.999	2
HAT-086-0004017	54.656399	79.24	0.014	0.004	1	-99.999999	-999.99	-9.999	-9.999	2
HAT-086-0004248	-99.999999	-999.99	-9.999	-9.999	2	-99.999999	-999.99	-9.999	-9.999	2
HAT-086-0005153	-99.999999	-999.99	-9.999	-9.999	2	-99.999999	-999.99	-9.999	-9.999	2
HAT-086-0005542	-99.999999	-999.99	-9.999	-9.999	2	-99.999999	-999.99	-9.999	-9.999	2
HAT-086-0005602	-99.999999	-999.99	-9.999	-9.999	2	42.385695	24.98	0.015	0.004	1

^a The first 1 (or 2) letter(s) in the subscript denote the period-finding algorithm (a: AoV, ah: AoVHarm, d: DACF), the last letter in the subscript denotes the light-curve type (e: EPD, t: TFA).^b The peak-to-peak amplitude. The first 2 (or 3) letter(s) in the subscript denote the period used, the last letter in the subscript denotes the light-curve type used to calculate the amplitude (e: EPD, t: TFA).^c Quality flag (0: robust detection, 1: questionable detection, 2: not detected by the given method for the given light curve type).

TABLE 9
CATALOG OF CANDIDATE PERIODIC VARIABLE STARS. III: AoVHARM

ID	P _{ahe} ^a d	SN _{ahe} ^a	Amp _{ahee} ^b mag	Amp _{ahet} ^b mag	Q _{ahe} ^c	P _{ah} ^a d	SN _{ah} ^a	Amp _{ah} ^b mag	Amp _{ah} ^b mag	Q _{ah} ^c
HAT-085-0004278	-99.999999	-999.99	-9.999	-9.999	2	-99.999999	-999.99	-9.999	-9.999	2
HAT-085-0053487	-99.999999	-999.99	-9.999	-9.999	2	-99.999999	-999.99	-9.999	-9.999	2
HAT-086-0001701	-99.999999	-999.99	-9.999	-9.999	2	17.887814	41.17	0.010	0.005	0
HAT-086-0002357	-99.999999	-999.99	-9.999	-9.999	2	-99.999999	-999.99	-9.999	-9.999	2
HAT-086-0003825	-99.999999	-999.99	-9.999	-9.999	2	-99.999999	-999.99	-9.999	-9.999	2
HAT-086-0004017	40.868328	95.12	0.014	0.011	1	-99.999999	-999.99	-9.999	-9.999	2
HAT-086-0004248	-99.999999	-999.99	-9.999	-9.999	2	-99.999999	-999.99	-9.999	-9.999	2
HAT-086-0005153	16.368924	49.27	0.010	0.003	0	-99.999999	-999.99	-9.999	-9.999	2
HAT-086-0005542	20.363047	64.94	0.010	0.009	0	20.411745	29.90	0.010	0.009	0
HAT-086-0005602	-99.999999	-999.99	-9.999	-9.999	2	21.071980	47.14	0.013	0.005	1

^a The first 1 (or 2) letter(s) in the subscript denote the period-finding algorithm (a: AoV, ah: AoVHarm, d: DACF), the last letter in the subscript denotes the light-curve type (e: EPD, t: TFA).^b The peak-to-peak amplitude. The first 2 (or 3) letter(s) in the subscript denote the period used, the last letter in the subscript denotes the light-curve type used to calculate the amplitude (e: EPD, t: TFA).^c Quality flag (0: robust detection, 1: questionable detection, 2: not detected by the given method for the given light curve type).

TABLE 10
CATALOG OF CANDIDATE PERIODIC VARIABLE STARS. IV: DACF EPD

ID	P_{de}^a d	$eP_{de}^{a,b}$ d	$y_{de}^{a,c}$	$Pr_{de}^{a,d}$	Amp_{dee}^e mag	Amp_{det}^e mag	$Pf_{de}^{a,f}$	$Q_{de}^{a,g}$
HAT-085-0004278	28.878214	0.744375	0.2040	-9.9122	0.021	0.006	0	1
HAT-085-0053487	-99.999999	-99.999999	-9.9999	999.9999	-9.999	-9.999	2	2
HAT-086-0001701	-99.999999	-99.999999	-9.9999	999.9999	-9.999	-9.999	2	2
HAT-086-0002357	28.562554	1.731345	0.0389	-21.1846	0.009	0.002	0	1
HAT-086-0003825	28.930191	0.703257	0.1421	-11.8251	0.009	0.002	1	0
HAT-086-0004017	40.432676	1.834407	0.0633	-12.3304	0.013	0.014	1	1
HAT-086-0004248	30.140252	0.840410	0.1182	-12.9545	0.011	0.002	1	1
HAT-086-0005153	50.034209	3.677269	0.1127	-13.5771	0.007	0.008	1	0
HAT-086-0005542	-99.999999	-99.999999	-9.9999	999.9999	-9.999	-9.999	2	2
HAT-086-0005602	64.435201	11.503159	0.0130	-11.8146	0.009	0.005	1	0

...

^a The first 1 (or 2) letter(s) in the subscript denote the period-finding algorithm (a: AoV, ah: AoVHarm, d: DACF), the last letter in the subscript denotes the light-curve type (e: EPD, t: TFA).^b Period uncertainty from fitting a quadratic function to the cross-correlation peak.^c Peak cross-correlation value^d Natural logarithm of the false-alarm probability^e The peak-to-peak amplitude. The first 2 (or 3) letter(s) in the subscript denote the period used, the last letter in the subscript denotes the light-curve type used to calculate the amplitude (e: EPD, t: TFA).^f Period flag (0: ok, 1: true period is likely a harmonic of measured period, 2: no detection).^g Quality flag (0: robust detection, 1: questionable detection, 2: not detected by the given method for the given light curve type).

TABLE 11
CATALOG OF CANDIDATE PERIODIC VARIABLE STARS. V: DACF TFA

ID	P_{dt}^a d	$eP_{dt}^{a,b}$ d	$y_{dt}^{a,c}$	$Pr_{dt}^{a,d}$	Amp_{dte}^e mag	Amp_{dtt}^e mag	$Pf_{dt}^{a,f}$	$Q_{dt}^{a,g}$
HAT-085-0004278	-99.999999	-99.999999	-9.9999	999.9999	-9.999	-9.999	2	2
HAT-085-0053487	-99.999999	-99.999999	-9.9999	999.9999	-9.999	-9.999	2	2
HAT-086-0001701	-99.999999	-99.999999	-9.9999	999.9999	-9.999	-9.999	2	2
HAT-086-0002357	-99.999999	-99.999999	-9.9999	999.9999	-9.999	-9.999	2	2
HAT-086-0003825	-99.999999	-99.999999	-9.9999	999.9999	-9.999	-9.999	2	2
HAT-086-0004017	-99.999999	-99.999999	-9.9999	999.9999	-9.999	-9.999	2	2
HAT-086-0004248	62.607406	36.891590	0.0150	-10.7230	0.014	0.004	1	1
HAT-086-0005153	-99.999999	-99.999999	-9.9999	999.9999	-9.999	-9.999	2	2
HAT-086-0005542	-99.999999	-99.999999	-9.9999	999.9999	-9.999	-9.999	2	2
HAT-086-0005602	20.668377	2.433455	0.0079	-16.9868	0.013	0.005	0	0

...

^a The first 1 (or 2) letter(s) in the subscript denote the period-finding algorithm (a: AoV, ah: AoVHarm, d: DACF), the last letter in the subscript denotes the light-curve type (e: EPD, t: TFA).^b Period uncertainty from fitting a quadratic function to the cross-correlation peak.^c Peak cross-correlation value^d Natural logarithm of the false-alarm probability^e The peak-to-peak amplitude. The first 2 (or 3) letter(s) in the subscript denote the period used, the last letter in the subscript denotes the light-curve type used to calculate the amplitude (e: EPD, t: TFA).^f Period flag (0: ok, 1: true period is likely a harmonic of measured period, 2: no detection).^g Quality flag (0: robust detection, 1: questionable detection, 2: not detected by the given method for the given light curve type).

TABLE 12
CATALOG OF CANDIDATE ECLIPSING BINARIES. I: ASTROMETRIC AND PHOTOMETRIC DATA

ID	RA ^a h:m:s	DE ^a deg:m:s	pmRA ^b mas/yr	pmDE ^b mas/yr	Jmag ^c mag	Hmag ^c mag	Kmag ^c mag	Vmag mag	Vref ^d	Bl ^e
HAT-086-0022206	23:50:17.08	+51:11:28.9	37.45	-20.48	11.655	11.009	10.858	13.656	1	1
HAT-087-0016301	00:34:50.23	+53:42:12.1	-37.41	12.76	11.310	10.613	10.433	13.904	2	2
HAT-088-0017192	01:38:56.50	+48:52:40.2	-11.17	-29.81	11.469	10.807	10.608	13.920	1	0
HAT-113-0005699	18:09:47.63	+49:02:55.0	-53.66	-21.85	11.362	10.710	10.560	14.126	1	0
HAT-127-0008153	03:04:05.20	+42:03:10.5	53.86	-54.80	10.650	9.983	9.730	14.159	1	2
HAT-127-0009895	03:02:22.68	+43:40:48.0	-76.01	-30.99	11.018	10.375	10.198	13.890	1	3
HAT-129-0025342	04:13:06.81	+41:21:01.4	42.67	-26.33	11.938	11.372	11.160	14.311	1	1
HAT-131-0026711	05:16:36.90	+48:35:44.3	-14.58	-32.73	11.802	11.138	10.982	14.275	1	2
HAT-132-0012475	05:43:20.18	+47:17:00.1	7.93	-29.15	11.345	10.704	10.569	13.765	1	2
HAT-133-0001901	06:31:57.29	+41:33:57.3	-56.65	-45.10	9.581	8.930	8.787	11.638	1	1

...

^a J2000, from 2MASS^b from PPMX^c from 2MASS^d Source for Vmag (0: PPMX (Tycho-2), 1: Transformed from USNO-B1.0, 2: Transformed from 2MASS)^e Blending Flag (0: unblended, 1: unlikely blend, 2: possible blend, 3: probable blend, 4: problematic)

TABLE 13
CATALOG OF CANDIDATE ECLIPSING BINARIES. II: EPHEMERIDES

ID	Per d	ePer d	HJD0p ^a d	eHJD0p ^b d	DURp ^c h	HJD0s ^a d	eHJD0s ^b d	DURs ^c h
HAT-086-0022206	0.4290339	0.0000062	2454431.11547	0.00042	1.53	2454436.90565	0.00117	1.52
HAT-087-0016301	3.1325961	0.0003763	2454427.60691	0.00504	5.24	2454416.64515	0.00442	4.63
HAT-088-0017192	1.4030920	0.0000252	2453693.67973	0.00066	2.15	2453680.35434	0.00071	2.00
HAT-113-0005699	0.2278737	0.0000013	2453901.88979	0.00028	1.24	2453905.42128	0.00039	1.14
HAT-127-0008153	2.5681924	0.0000869	2452979.53695	0.00140	1.98	2452991.10638	0.00215	1.53
HAT-127-0009895	0.5916726	0.0000239	2452985.85734	0.00187	4.16	2452990.88691	0.05334	4.55
HAT-129-0025342	4.6109145	0.0002340	2453726.54374	0.00129	2.85	2453724.34418	0.00235	3.34
HAT-131-0026711	0.6639531	0.0000048	2454497.17095	0.00026	1.43	2454498.83198	0.00170	1.42
HAT-132-0012475	0.8663223	0.0000035	2454128.11503	0.00145	3.12	9999999.99999	9.99999	9.99
HAT-133-0001901	0.2929328	0.0000001	2453390.29599	0.00016	0.91	2453376.38208	0.00022	0.88

...

^a Epoch of primary (p) or secondary (s) eclipse. The value is 9999999.99999 if a secondary eclipse is not detected.^b Uncertainty on the epoch of eclipse.^c Eclipse duration.

TABLE 14
CATALOG OF FLARE EVENTS. I: ASTROMETRIC AND PHOTOMETRIC DATA^a

ID	RA ^b h:m:s	DE ^b deg:m:s	pmRA ^c mas/yr	pmDE ^c mas/yr	Jmag ^d mag	Hmag ^d mag	Kmag ^d mag	Vmag mag	Vref ^e
HAT-087-0002617	00:21:57.81	+49:12:38.0	205.00	-32.56	9.139	8.454	8.205	12.474	1
HAT-094-0007885	05:35:22.84	+53:16:02.7	4.39	-35.34	10.797	10.139	9.958	13.797	1
HAT-101-0000851	09:59:39.19	+48:47:54.6	-45.67	8.38	9.686	9.002	8.817	12.137	1
HAT-103-0001602	11:03:17.04	+49:48:25.5	-63.32	-7.74	10.898	10.286	10.114	13.544	1
HAT-119-0031226	22:02:08.40	+49:07:17.3	-68.41	0.39	11.102	10.434	10.228	12.777	1
HAT-123-0002064	00:53:06.48	+48:29:38.5	220.82	-162.32	9.427	8.773	8.580	12.899	0
HAT-123-0002064	00:53:06.48	+48:29:38.5	220.82	-162.32	9.427	8.773	8.580	12.899	0
HAT-123-0003289	00:41:41.41	+44:10:53.1	-46.23	-15.14	10.012	9.399	9.188	14.251	1
HAT-126-0002065	02:40:52.51	+44:52:36.5	215.74	3.85	9.277	8.696	8.461	13.197	0
HAT-128-0007422	03:42:45.17	+43:27:40.1	30.96	-64.10	10.435	9.830	9.568	14.611	1

...

^a Stars may be listed more than once if they have multiple flare events^b J2000, from 2MASS^c from PPMX^d from 2MASS^e Source for Vmag (0: PPMX (Tycho-2), 1: Transformed from USNO-B1.0, 2: Transformed from 2MASS)

TABLE 15
CATALOG OF FLARE EVENTS. II: FLARE DATA^a

ID	HJD ^b d	Amp ^c	τ ^d d	Nobs ^e
HAT-087-0002617	2454437.61429	0.130	0.0047	26
HAT-094-0007885	2454463.88452	0.219	0.0092	7
HAT-101-0000851	2454162.63179	0.200	0.0176	26
HAT-103-0001602	2454563.96193	0.279	0.0212	29
HAT-119-0031226	2453360.74326	0.552	0.0346	31
HAT-123-0002064	2454337.05672	0.118	0.0053	11
HAT-123-0002064	2454391.64587	0.482	0.0065	15
HAT-123-0003289	2454338.90758	0.236	0.0087	23
HAT-126-0002065	2452999.82533	0.277	0.0056	9
HAT-128-0007422	2453387.68872	0.346	0.0216	13
...				

^a Stars may be listed more than once if they have multiple flare events^b Time of flare peak^c Peak intensity relative to the non-flaring intensity.^d 1/e decay time of the flare.^e Number of observations included in fitting the flare.

TABLE 16
CROSS-MATCHES TO PREVIOUSLY IDENTIFIED VARIABLE STARS

ID	Alternate ID	Mflag ^a	BlFlag ^b
HAT-086-0005602	V0544 Cas	1	3
HAT-086-0022206	V1001 Cas	0	1
HAT-091-0010976	V0524 Per	0	2
HAT-091-0016338	V0485 Per	0	1
HAT-112-0001188	1SWASP J173659.28+485946.1	0	0
HAT-113-0005699	ROTSE1 J180947.50+490254.5	0	0
HAT-113-0005699	V1104 Her	0	0
HAT-121-0003519	NSV 14597	0	3
HAT-128-0006973	NSV 01380	1	3
HAT-128-0015056	NSVS 15796	1	3
...			

^a 0: the match is correct, 1: the two stars are not the same, but may be blended in the HAT images.^b Automatically generated blending flag. 0: unblended, 1: unlikely blend, 2: potential blend, 3: probable blend, 4: problematic, 5: source is a flare star, no blending info is generated.

TABLE 17
CROSS-MATCHES TO ROSAT X-RAY SOURCES

ID	RosatID ^a	lgfxfj ^b
HAT-087-0001873	1RXS J002854.1+502226	-1.7955
HAT-087-0002547	1RXS J003633.7+553728	-2.2528
HAT-087-0002617	1RXS J002158.0+491245	-2.1767
HAT-093-0001111	1RXS J044853.2+552725	-2.3278
HAT-094-0004894	1RXS J050718.8+530713	-2.3805
HAT-094-0005389	1RXS J053001.7+484932	-2.0147
HAT-094-0006428	1RXS J050809.3+532437	-1.7462
HAT-094-0009460	1RXS J050012.4+532033	-1.6652
HAT-094-0384094	1RXS J053625.1+533618	-2.2575
HAT-103-0001602	1RXS J110318.0+494822	-1.8031
...		

^a ID of matching ROSAT source^b Estimated $\log(f_x/f_J)$ (ratio of X-ray flux to J-band flux).